

## **Product datasheet for CF813957**

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# FABP3 Mouse Monoclonal Antibody [Clone ID: OTI2B12]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2B12

**Applications:** ELISA

Recommended Dilution: ELISA 1:5000-10000

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human FABP3 (NP\_004093) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Predicted Protein Size:** 14.9 kDa

**Gene Name:** fatty acid binding protein 3

Database Link: NP 004093

Entrez Gene 2170 Human

P05413



Synonyms:

#### FABP3 Mouse Monoclonal Antibody [Clone ID: OTI2B12] - CF813957

**Background:** The intracellular fatty acid-binding proteins (FABPs) belongs to a multigene family. FABPs are

divided into at least three distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15 kDa proteins and are thought to participate in the uptake, intracellular metabolism and/or transport of long-chain fatty acids. They may also be responsible in the modulation of cell growth and proliferation. Fatty acid-binding protein 3 gene contains four exons and its function is to arrest growth of mammary epithelial cells. This gene is a candidate tumor suppressor gene for human breast cancer. Alternative splicing results in

multiple transcript variants. [provided by RefSeq, Mar 2016]

FABP11; H-FABP; M-FABP; MDGI; O-FABP

**Protein Pathways:** PPAR signaling pathway