

Product datasheet for CF814597

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL17 (IL17A) Mouse Monoclonal Antibody [Clone ID: OTI7H6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7H6

Applications: ELISA

Recommended Dilution: ELISA 1:5000-10000

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 24-155 of human IL-17A

(NP_002181) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Predicted Protein Size: 17.5 kDa

Gene Name: interleukin 17A

Database Link: NP 002181

Entrez Gene 3605 Human

Q16552





IL17 (IL17A) Mouse Monoclonal Antibody [Clone ID: OTI7H6] - CF814597

Background: The protein encoded by this gene is a proinflammatory cytokine produced by activated T

cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis,

psoriasis and multiple sclerosis. [provided by RefSeq, Jul 2008]

Synonyms: CTLA-8; CTLA8; IL-17; IL-17A; IL17; ILA17

Protein Families: Druggable Genome, Secreted Protein
Protein Pathways: Cytokine-cytokine receptor interaction