

Product datasheet for **CL005P**

Cd5 (Ly-1.1) Mouse Monoclonal Antibody [Clone ID: 7-20.6/3]

Product data:

Product Type:	Primary Antibodies
Clone Name:	7-20.6/3
Applications:	CT, FC
Recommended Dilution:	Cytotoxicity tests. Flow Cytometry.
Reactivity:	Mouse
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Recipient: 129/ReJ. Donor: B6-Ly-1a. Fusion Partner: Spleen from immunized recipient fused with myeloma P3-NSI-Ag4-1.
Specificity:	Anti-Ly 1.1 monoclonal antibody reacts with T lymphocytes from mouse strains expressing the Ly 1.1 phenotype, but does not react with lymphocytes from mouse strains expressing the Ly 1.2 phenotype. This antibody can be used to identify T helper cells as they have a high density of Ly 1 on their surface.
Formulation:	PBS without preservatives State: Purified State: Liquid (0.2 µm filtered) purified Ig fraction
Concentration:	lot specific
Purification:	Protein G Chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	CD5 antigen



[View online »](#)

Database Link: [Entrez Gene 12507 Mouse P13379](#)

Background: CD5 is a 55kDa T lymphocyte single chain transmembrane glycoprotein. It is present on all mature T lymphocytes, on most thymocytes and on many T cell leukemias and lymphomas. It reacts with a subpopulation of activated B cells. CD5/Lyt1 antigen is a monomeric type I transmembrane glycoprotein expressed on thymocytes, T lymphocytes, and a subset of B lymphocytes, but not on natural killer (NK) cells. It has been identified as the major ligand of the B cell antigen CD72. The frequency of CD5+ B cells exhibits strain dependent variation, and the phenotypic, anatomical, functional, developmental, and pathological characteristics of the CD5+ B cells suggest that they may represent a distinct lineage, known as B1 cells. Binding of CD5 on the T cell surface can augment alloantigen or mitogen induced lymphocyte proliferation and induces increased cytosolic free calcium, IL2 secretion, and IL2R expression. It has been proposed that CD5 negatively regulates signal transduction mediated by the T cell and B cell receptors.

Synonyms: CD5, LEU1

Note: Protocol: **Flow Cytometry Analysis:**

Method:

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte®-M cell separation medium.
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in media A. Add 50 μ l of this suspension to each tube (each tube will then contain 1×10^6 cells, representing 1 test).
4. To each tube, add 0.2-0.5 μ g* of CL005P.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Add 100 μ l of FITC Goat anti-Mouse IgG (H+L) secondary antibody at a 1:500 dilution.
9. Incubate the tubes at 4°C for 30-60 minutes.
(It is recommended that the tubes are protected from light since most fluorochromes are light sensitive).
10. Wash 2 times at 4°C in media B.
11. Resuspend the cell pellet in 50 μ l ice cold media B.
12. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 μ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 μ l of 2M sodium azide in 100 mls).

Results:
Tissue Distribution by Flow Cytometry Analysis:

Mouse Strain: C3H/He

Cell Concentration : 1x10⁶ cells per tests

Antibody Concentration Used: 0.5 µg/10⁶ cells

Isotypic Control: Purified Mouse IgG2a

Cytotoxicity Analysis:
Method:

1. Prepare a cell suspension from the appropriate tissue in Cytotoxicity Medium or equivalent. Remove red cells and dead cells (where necessary) by purification of viable lymphocytes on Lympholyte®-M (b) density cell separation medium. After washing, adjust the cell concentration to 1x10⁶ cells per ml in Cytotoxicity Medium.
 2. Add the antibody to a final concentration of 1:500 and mix.
 3. Incubate for 60 minutes at 4°C.
 4. Centrifuge to pellet the cells and discard the supernatant.
 5. Resuspend to the original volume in Low-Tox®-M Rabbit Complementc diluted 1:12 in Cytotoxicity Medium.
 6. Incubate for 60 minutes at 37°C.
 7. Place on ice.
 8. Add Trypan Blue, 10% by volume of 1% Trypan Blue (w/v) added 3-5 minutes before scoring works well. Score live versus dead cells in a hemacytometer.
- C.I. = Cytotoxicity Index = 100x

$$\frac{\% \text{Cyt}(\text{Ab} + \text{Complement}) - \% \text{Cyt}(\text{Complement})}{100\% - \% \text{Cyt}(\text{Complement})}$$

Notes:

- a. Cytotoxicity Medium is RPMI-1640 with 25 mM Hepes buffer and 0.3% bovine serum albumin (BSA). BSA is substituted for the conventionally used fetal calf serum (FCS) because we have found that many batches of FCS contain complement dependent cytotoxins to mouse lymphocytes, thus increasing the background killing in the presence of complement. We recommend that cells not be exposed to FCS prior to or during exposure to antibody and complement. Some batches of BSA also contain complement dependent cytotoxins, resulting in the same problem. We screen for batches of BSA giving low background in the presence of complement and use the selected BSA for preparing Cytotoxicity Medium.
- b. Lympholyte®-M cell separation medium is density separation medium designed specifically for the isolation of viable mouse lymphocytes. This separation medium provides a high and non-selective recovery of viable mouse lymphocytes, removing red cells and dead cells. The density of this medium is 1.087 - 1.088. Isolation of mouse lymphocytes on cell separation medium of density 1.077 will result in high and selective loss of lymphocytes and should be avoided.
- c. Rabbit serum provides the most potent source of complement for use with antibodies to mouse cell surface antigens. However, rabbit serum itself is very toxic to mouse lymphocytes. Low-Tox®-M Rabbit Complement is absorbed to remove toxicity to mouse lymphocytes, while maintaining its high complement activity. When used in conjunction with Cytotoxicity

Medium, this reagent provides a highly potent source of complement with minimal background toxicity.

Results:

Antibody Titration by Cytotoxicity Analysis:

Cell Source: Thymocytes

Donor: C3H/He

Cell Concentration: 1.1x10⁶ cells/ml

Complement: Low-Tox®-M Rabbit Complement

Complement Concentration: 1:15

C.I. = Cytotoxicity Index = 100x

%Cyt(Ab+Complement) - %Cyt (Complement)/100%- %Cyt (Complement)

Tissue Distribution by Cytotoxicity Analysis:

Antibody Concentration: 1:640

Strain: C3H/He

Cell Source (C.I.)

Thymus (99)

Spleen (28)

Lymph Node (71)

Strain Distribution by Cytotoxicity Analysis:

Antibody Concentration Used: 1: 500

Strains Tested: BALB/c, A.TH, CBA/J, C3H/He, C57BL/6

Positive: CBA/J, C3H/He

Negative: C57BL/6, BALB/c, A.TH

Functional Analysis:

Method:

Cells were treated as described in **Cytotoxic Depletion of CD5 (Ly 1.2) Positive Lymphocytes**. Treated cells and controls were tested for:

- the ability to generate plaque-forming cells (PFC) using a modified Jerne haemolytic plaque assay
- the ability to generate cytotoxic T effector cells using a cytotoxic lymphocyte reaction (CTL) assay.

Cells were treated both before and after sensitization in the CTL assay. In vitro immunizations were used in all experiments.

Results:

Cell Source: Splenocytes

Donors: C57BL/6 and C3H/He

Cell Concentration: 1x10⁷ cells/ml

Antibody Concentration used: 1:10

Complement: Low-Tox®-M Rabbit Complement


Complement Concentration: 1:10

Treatment of C3H/He splenocytes with anti-CD5 (Ly 1.1) plus complement was found to reduce the number of plaque-forming cells and inhibit cytotoxic T cell generation. Cytotoxic T effector cell function was not affected (cells treated after sensitization). No effect was observed when C57BL/6 cells were used. These results are consistent with the depletion of T helper cells of the Ly 1.1 phenotype.

Cytotoxic Depleting Of CD5 (Ly 1.1) Positive Lymphocytes:

1. Prepare a cell suspension from the appropriate tissue in Cytotoxicity Medium (a) or equivalent. Remove red cells and dead cells (where necessary) by purification of viable lymphocytes on Lympholyte®-M cell separation medium. After washing, adjust the cell concentration to 1×10^7 cells per ml in Cytotoxicity Medium.
2. Add the antibody to a final concentration of 1:500 and mix. Alternatively, pellet the cells and resuspend in antibody diluted 1:500 in Cytotoxicity Medium.
3. Incubate for 60 minutes at 4°C.
4. Centrifuge to pellet the cells and discard the supernatant.
5. Resuspend to the original volume in Low-Tox-M® Rabbit Complement, diluted to the appropriate concentration in Cytotoxicity Medium. (Recommended concentration included with each batch of Low-Tox-M® Rabbit Complement.)
6. Incubate for 60 minutes at 37°C.
7. Monitor for percent cytotoxicity at this stage, before further processing. For this purpose, remove a small sample from each tube, dilute 1:10 with medium, and add 1/10 volume of 1% Trypan Blue. After 3-5 minutes, score live versus dead cells in a hemacytometer.
8. For functional studies, remove the dead cells from the treated groups before further processing, particularly if the treated cells are to be cultured. This can be done by layering the cell suspension over a separation medium and centrifuging at room temperature as per the instructions provided. Live cells will form a layer at the interface, while the dead cells pellet. The interface can then be collected and washed in Cytotoxicity Medium before being resuspended in the appropriate medium for further processing. Alternatively, the cells can be washed and resuspended in the appropriate medium for further processing immediately after Step #6, provided that the dead cells will not interfere with subsequent assays.

Product images:



Cell Source: Thymus. Percentage of cells stained above control: 97.9%

