

## Product datasheet for **CL029F**

### **CD45 / LCA (CD45RB) Rat Monoclonal Antibody [Clone ID: 16A]**

#### **Product data:**

<b>Product Type:</b>	Primary Antibodies
<b>Clone Name:</b>	16A
<b>Applications:</b>	FC
<b>Recommended Dilution:</b>	Flow cytometry. Immunohistochemistry on acetone-fixed frozen sections and paraffin embedded sections. Immunoprecipitation.
<b>Reactivity:</b>	Mouse
<b>Host:</b>	Rat
<b>Isotype:</b>	IgG2a
<b>Clonality:</b>	Monoclonal
<b>Immunogen:</b>	TH2 cell clones, final boost with TH2 clone D10 Fusion Partner: Ag8.653
<b>Specificity:</b>	This monoclonal antibody reacts with the CD45 isoform containing the exon B dependent epitope. CD45RB is highly expressed on peripheral B cells, cytotoxic T cells, a subset of T helper cells and most thymocytes.
<b>Formulation:</b>	PBS, 0.09% NaN <sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. Label: FITC State: Liquid
<b>Concentration:</b>	lot specific
<b>Purification:</b>	Protein G Chromatography
<b>Conjugation:</b>	FITC
<b>Storage:</b>	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. This product is photosensitive and should be protected from light.
<b>Stability:</b>	Shelf life: one year from despatch.
<b>Gene Name:</b>	protein tyrosine phosphatase, receptor type, C



[View online »](#)

**Database Link:** [Entrez Gene 19264 Mouse P06800](#)

**Background:** CD45 is a family of single chain transmembrane glycoproteins consisting of at least four isoforms (220, 205, 190, 180 kDa) which share a common large intracellular domain. Their extracellular domains are heavily glycosylated. The different isoforms are produced by alternative messenger RNA splicing of three exons of a single gene on chromosome 1. CD45 is expressed on cells of the human hematopoietic lineage (including hematopoietic stem cells) with the exception of mature red cells. It is not detected on differentiated cells of other tissues. It is likely that CD45 plays an important role in signal transduction, inhibition or upregulation of various immunological functions. Antibodies recognising a common epitope on all of the isoforms are termed CD45 whilst those recognising only individual isoforms are termed CD45RA or CD45RO etc.

**Synonyms:** PTPRC, Leukocyte common antigen, L-CA, T200

**Note:** Protocol: **FLOW CYTOMETRY ANALYSIS:**

**Method:**

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte®-M cell separation medium.
2. Wash 2 times.
3. Resuspend the cells to a concentration of  $2 \times 10^7$  cells/ml in media A. Add 50  $\mu$ l of this suspension to each tube (each tube will then contain  $1 \times 10^6$  cells, representing 1 test).
4. To each tube, add  $\sim 1.0 \mu\text{g}^*$  of this Ab per  $10^6$  cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at  $4^\circ\text{C}$ . (It is recommended that the tubes are protected from light, since most fluorochromes are light sensitive.)
7. Wash 2 times at  $4^\circ\text{C}$ .
8. Resuspend the cell pellet in 50  $\mu$ l ice cold media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15  $\mu$ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

**Media:**

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100  $\mu$ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100  $\mu$ l of 2M sodium azide in 100 mls).

**Results - Tissue Distribution:**


Mouse Strain: BALB/c

Cell Concentration:  $1 \times 10^6$  cells per test

Antibody Dilution Used: 1.0  $\mu\text{g}/10^6$  cells

Isotypic Control: FITC Rat IgG2a

## Product images:



Cell Source: Spleen

Percentage of cells stained above control: 79.8%