

## Product datasheet for **CL032A**

### Sell Rat Monoclonal Antibody [Clone ID: MEL-14]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	MEL-14
Applications:	FC, FN, IHC, IP
Recommended Dilution:	Flow cytometry. Immunohistochemistry. Immunoprecipitation.
Reactivity:	Mouse
Host:	Rat
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Mouse B cell Lymphoma, 38C-14 Donor: Fischer Rat Spleen Fusion Partner: P3 X 63Ag8.653
Specificity:	This monoclonal antibody reacts with a 90 kDa protein which is involved with the homing of lymphocytes to peripheral lymph nodes.
Formulation:	PBS, no preservative, 0.2 µm filtered State: Azide Free State: Liquid
Concentration:	lot specific
Purification:	Protein G Chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	selectin, lymphocyte
Database Link:	<a href="#">Entrez Gene 20343 Mouse P18337</a>



[View online »](#)

**Background:** L-selectin is expressed on most T and B lymphocytes, neutrophils, monocytes, eosinophils.<sup>1</sup> Pre-incubation of lymphocytes with this antibody completely and specifically blocks binding of lymphocytes to high endothelial venules (HEV) in vitro<sup>2,3,6</sup> and the migration of lymphocytes to lymph nodes in vivo.<sup>2,3</sup> Polymorphonuclear cells preincubated with this antibody do not migrate to the inflammatory foci.

**Synonyms:** SELL, LNHR, LYAM1, Leu-8, TQ1, gp90-MEL, LECAM1, LAM-1

**Note:** Protocol: **FLOW CYTOMETRY ANALYSIS:**

**Method:**

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte®-M cell separation medium.
2. Wash 2 times.
3. Resuspend the cells to a concentration of  $2 \times 10^7$  cells/ml in media A. Add 50  $\mu$ l of this suspension to each tube (each tube will then contain  $1 \times 10^6$  cells, representing 1 test).
4. To each tube, add 0.1-0.2  $\mu$ g\* of this Ab.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Add 100  $\mu$ l of secondary antibody (FITC Goat anti-rat IgG (H+L)) at 1: 500 dilution.
9. Incubate the tubes at 4°C for 30-60 minutes. (It is recommended that the tubes are protected from light since most fluorochromes are light sensitive).
10. Wash 2 times at 4°C in media B.
11. Resuspend the cell pellet in 50  $\mu$ l ice cold media B.
12. Transfer to suitable tubes for flow cytometric analysis containing 15  $\mu$ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

**Media:**

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100  $\mu$ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100  $\mu$ l of 2M sodium azide in 100 mls).

**Results - Tissue Distribution:**

Mouse Strain: BALB/c

Cell Concentration:  $1 \times 10^6$  cells per test

Antibody Concentration Used: 0.2  $\mu$ g/ $10^6$  cells

Isotypic Control: Rat IgG2a

**Cell Source: Percentage of cells stained above control**

Thymus: 85.8%

Spleen: 41.0%

Lymph Node: 75.0%

**Results - Strain Distribution:**

Tissue: Spleen

Cell Concentration: 1x10e6 cells per test


Antibody Concentration Used: 0.2 µg/10e6 cells

Strains Tested: BALB/c, C57BL/6, C3H/He, CBA/J, AKR

Positive: BALB/c, C57BL/6, C3H/He, CBA/J, AKR

Negative: none

**Product images:**



Cell Source: Lymph Node

Percentage of cells stained above control: 75.0%