

Product datasheet for **CL042RX**

Kit Rat Monoclonal Antibody [Clone ID: ACK4]

Product data:

Product Type:	Primary Antibodies
Clone Name:	ACK4
Applications:	FC
Recommended Dilution:	Flow Cytometry.
Reactivity:	Mouse
Host:	Rat
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	IL-3 dependent mast cells derived from WB- +/- mice Donor: Wistar spleen Fusion Partner: X63.653. Ag8
Specificity:	Anti-mouse CD117 monoclonal antibody recognizes the receptor tyrosine kinase, c-kit. The ligand for this receptor is steel factor (stem cell factor), which exists in both soluble and membrane form. The interaction between steel factor and c-kit is essential for the development of hematopoietic, gonadal and pigment stem cells. c-kit positive cells are a subset of CD34+ hematopoietic precursor cells and it is expressed on 5-10% of total adult bone marrow cells.
Formulation:	PBS containing 0.02% sodium azide (NaN ₃) as preservative and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. Label: PE State: Liquid purified Ig fraction Label: R - Phycoerythrin
Concentration:	lot specific
Purification:	Affinity chromatography on Protein G
Conjugation:	PE
Storage:	Store the antibody undiluted at 2-8°C. DO NOT FREEZE!
Stability:	Shelf life: one year from despatch.



[View online »](#)

Gene Name:	kit oncogene
Database Link:	Entrez Gene 16590 Mouse P05532
Synonyms:	SCFR, KIT
Note:	<p>Protocol: <u>FLOW CYTOMETRY ANALYSIS:</u></p> <p>Method:</p> <ol style="list-style-type: none">1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte®-M (cell separation medium).2. Wash 2 times.3. Resuspend the cells to a concentration of 2x10⁷ cells/ml in media A. Add 50 µl of this suspension to each tube (each tube will then contain 1 x 10⁶ cells, representing 1 test).4. To each tube, add 0.5 – 1.0 µg of this antibody per 10⁶ cells.5. Vortex the tubes to ensure thorough mixing of antibody and cells.6. Incubate the tubes for 30 minutes at 4°C. (It is recommended that the tubes are protected from light, since most fluorochromes are light sensitive.)7. Wash 2 times at 4°C.8. Resuspend the cell pellet in 50 µl ice cold media B.9. Transfer to suitable tubes for flow cytometric analysis containing 15 µl of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA. <p>Media:</p> <p>A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 µl of 2M sodium azide in 100 mls).</p> <p>B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 µl of 2M sodium azide in 100 mls).</p> <p>Results:</p> <p><u>Tissue Distribution by Flow Cytometry Analysis:</u></p> <p>Mouse Strain: BALB/c Cell Concentration: 1x10⁶ cells per test Antibody Concentration Used: 0.5 µg/10⁶ cells Isotypic Control: PE Rat IgG2a</p> <p><u>Strain Distribution by Flow Cytometry Analysis:</u></p> <p>Cell Concentration: 1x10⁶ cells per test Antibody Concentration Used: 1.0 mg /10⁶ cells Strains Tested: AKR, BALB/c, C3H/He, C57BL/6 Positive: AKR, BALB/c, C3H/He, C57BL/6 Negative: none</p>

Product images:

