

Product datasheet for CL044R1

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Tnfrsf1a Rat Monoclonal Antibody [Clone ID: HM104]

Product data:

Product Type: Primary Antibodies

Clone Name: HM104

Applications: FC

Recommended Dilution: Flow Cytometry: Use 10 μl of neat antibody to label 1% cells in 100 μl.

Reactivity: Mouse

Host: Rat

Isotype: IgG2a

Clonality: Monoclonal

Specificity: This antibody recognizes an extracellular region of Mouse CD120a (TNFR1, p55). Other

species not tested.

Formulation: PBS

Label: PE

State: Liquid purified IgG fraction

Stabilizer: 0.5% BSA

Preservative: 0.02% Sodium Azide Label: R. Phycoerythrin (RPE)

Purification: Ion Exchange Chromatography

Conjugation: PE

Storage: Store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Stability: Shelf life: One year from despatch.

Gene Name: tumor necrosis factor receptor superfamily, member 1a

Database Link: Entrez Gene 21937 Mouse

P25118





Tnfrsf1a Rat Monoclonal Antibody [Clone ID: HM104] - CL044R1

Background:

TNF-RI (~55-60 kDa) is present on most cell types and is considered to play a prominent role in cell stimulation by TNF-alpha. TNF-alpha activates inflammatory responses, induces apoptosis, regulates cellular proliferation, and may even promote cancer progression. The effects of TNF-alpha are mediated by TNF-RI and TNF-RII, which have both distinct and overlapping downstream signaling cascades. Induction of cytotoxicity and other functions are mediated largely via TNF-RI. TNF-RI is equally well activated by both the 17 kDa soluble and 26 kDa membrane-bound form, whereas TNF-RII is efficiently activated only by the membrane bound form of TNF-alpha. TNF-RI signaling is initiated when trimeric TNF-alpha binds TNF-RI receptors. Subsequent TNF-RI trimerization promotes the recruitment of a proximal signaling complex composed of TNF Receptor Associated protein with a Death Domain (TRADD), Receptor Interacting Protein (RIP), cellular Inhibitor of Apoptosis Protein 1 (cIAP1), TNF Receptor Associated Factor 2 (TRAF2), and likely TRAF5. Studies with TNF-RI-deficient mice indicate that TNF-RI mediates most of the proliferation, pro-inflammatory, and apoptosis-activating pathways.

Synonyms:

Tumor necrosis factor receptor 1, TNF-R1, TNF-RI, TNFR-I, p55, p60, Tnfrsf1a