

Product datasheet for **CL061**

H2-Q6 Mouse Monoclonal Antibody [Clone ID: 141-16.6]

Product data:

Product Type:	Primary Antibodies
Clone Name:	141-16.6
Applications:	CT, FC
Recommended Dilution:	Cytotoxicity Analysis. Flow Cytometry.
Reactivity:	Mouse
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Recipient: C58/J Donor: BALB/c-H-2b Fusion Partner: P3-NSI/1-Ag4(NS-1)
Specificity:	This monoclonal antibody detects the Qa.m2 antigenic determinant on the Qa-2 molecule. Thus, although the role of the Qa-2 antigen is unclear, antibodies to this antigen may be useful tools for separating functionally distinct subpopulations of lymphocytes and resolving pathways of differentiation. It may prove to be a useful marker in the study of B lymphocyte differentiation as well as differentiation of certain T cell functional populations.
Formulation:	State: Ascites State: Lyophilized ascites
Reconstitution Method:	Restore with 0.5 ml of distilled water.
Conjugation:	Unconjugated
Storage:	Prior to and following reconstitution store the antibody at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	histocompatibility 2, Q region locus 6
Database Link:	Entrez Gene 110557 Mouse P79568



[View online »](#)

Background:

The Qa region, so named because of the closely linked Qa 1-9 loci, is situated on murine chromosome 17 between the H-2D and T1a loci. These Qa loci code for cell surface alloantigens expressed on some lymphoid and myeloid cells. Their expression appears to be controlled by a gene or genes closely linked to the H-2D locus and they have been identified as being structurally related to the murine H-2 antigens. Although no functional role for Qa antigens has been determined, the restricted and differential distribution of Qa determinants on hemopoietic and lymphoid cells raises the possibility that they may not be simply differentiation markers but may play a role in hematopoietic differentiation.

The Qa-2 molecule has been identified as a two-chain structure composed of a 39,000 d heavy chain and a 12,000 d light chain which is B2 microglobulin. (The heavy chain is readily distinguished from that of H-2.) The Qa.m2 antigen has a wide representation on hematopoietic cells. It is however, found predominantly on peripheral T cells (cytotoxic and helper T cells) but was present on only 10% of thymus cells. Subpopulations of B cells (approximately 25% of all B cells) and bone marrow cells (15-20%) are also reactive. Qa.m2 was also detected on megakaryocyte colony-forming cells (MEG-CFC) but not on granulocyte-macrophage, erythroid or eosinophil colonies. In addition, the Qa.m2 is a marker of T cell differentiation, as it is selectively expressed on functional T cells. The control of Qa-2 antigen expression and of other Qa antigens by the H-2 region may reflect a control mechanism of functional T cell differentiation and ontogeny in developing T lymphocytes.

Note:

- a) The antibody has an identical strain distribution pattern to the Qa-2 conventional antisera, but shows some differences in tissue distribution.
- b) H-2Db strains carry approximately 8-10 times greater amounts of Qa-2 than strains carrying the H-2Dd or H-2Dq strains.

The reagent is not sold as sterile, but can be sterilized by filtration if necessary. To minimize loss of volume during filtration, dilute to the final working concentration in the appropriate medium before filtration and filter through a 0.45 μ millipore filter (or equivalent).

Protocol: RECOMMENDED METHOD FOR DEPLETING A CELL POPULATION OF Qa.m2 POSITIVE LYMPHOCYTES

1. Prepare a cell suspension from the appropriate tissue in Cytotoxicity Medium or equivalent. Remove red cells and dead cells (where necessary) by purification of viable lymphocytes on Lympholyte®-M density cell separation medium. After washing adjust the cell concentration to 1×10^7 cells per ml in Cytotoxicity Medium.
2. Add the antibody to a final concentration 1:20 and mix. Alternately, pellet the cells and resuspend in antibody diluted 1:20 in Cytotoxicity Medium.
3. Incubate for 60 minutes at 4°C.
4. Centrifuge to pellet the cells and discard the supernatant.
5. Resuspend to the original volume in Low-Tox®-M Rabbit Complement3, diluted to the appropriate concentration in Cytotoxicity Medium. (Recommended concentration included with each batch of Low-Tox®-M Rabbit Complement.)
6. Incubate for 60 minutes at 37°C.

7. Monitor for percent cytotoxicity at this stage, before further processing. For this purpose, remove a small sample from each tube, dilute 1:10 with medium, and add 1/10 volume of 1% trypan blue. After 3-5 minutes, score live vs. dead cells in a hemacytometer.
8. For functional studies, remove the dead cells from the treated groups before further processing, particularly if the treated cells are to be cultured. This can be done by layering the treated cell suspensions over an equal volume of Lympholyte®-M cell separation medium and centrifuging at room temperature as per the instruction provided. Live cells will form a layer at the interface, while the dead cells pellet. The interface can then be collected and washed in Cytotoxicity Medium before being resuspended in the appropriate medium for further processing. Alternately, the cells can be washed and resuspended in the appropriate medium for further processing immediately after Step #6, provided that the dead cells will not interfere with subsequent assays.

RECOMMENDED METHOD FOR DETERMINING PERCENT OF Qa.m2 POSITIVE CELLS IN A POPULATION

1. Prepare a cell suspension from the appropriate tissue in Cytotoxicity Medium1 or equivalent. Remove red cells and dead cells (where necessary) by purification of viable lymphocytes on Lympholyte®-M density cell separation medium. After washing, adjust the cell concentration to 1×10^6 cells per ml in Cytotoxicity Medium.
2. Add the antibody to a final concentration of 1:40 and mix.
3. Incubate for 60 minutes at 4°C.
4. Centrifuge to pellet the cells and discard the supernatant.
5. Resuspend to the original volume in Low-Tox®-M Rabbit Complement3, diluted to the appropriate concentration in Cytotoxicity Medium. (Recommended concentration included with each batch of Low-Tox®-M Complement).
6. Incubate for 60 minutes at 37°C.
7. Place on ice.
8. Add trypan blue. 10% volume of 1% trypan blue (w/v) added 3-5 minutes before scoring works well. Score live vs. dead cells in a hemacytometer. Cytotoxic Index (C.I.) can be calculated as shown in **FIGURE 1**.

Notes:

1. Cytotoxicity Medium® is RPMI-1640 with 25mM Hepes buffer and 0.3% bovine serum albumin (BSA). BSA is substituted for the conventionally used fetal calf serum (FCS) because we have found that many batches of FCS contain complement dependent cytotoxins to mouse lymphocytes, thus increasing the background killing in the presence of complement. We recommend that cells not be exposed to FCS prior to or during exposure to antibody and complement. Some batches of BSA also contain complement dependent cytotoxins, resulting in the same problem. We screen for batches of BSA giving low background in the presence of complement and use the selected BSA for preparing Cytotoxicity Medium®.
2. Lympholyte®-M cell separation medium is a density separation medium designed specifically for the isolation of viable mouse lymphocytes. This separation medium provides a high and non-selective recovery of viable mouse lymphocytes, removing red cells and dead cells. The density of this medium is 1.087-1.088. Isolation of mouse lymphocytes on cell

separation medium of density 1.077 will result in high and selective loss of lymphocytes and should be avoided.

3. Rabbit serum provides the most potent source of complement for use with antibodies to mouse cell surface antigens. However, rabbit serum itself is very toxic to mouse lymphocytes. Low-Tox®-M Rabbit Complement is absorbed to remove toxicity to mouse lymphocytes, while maintaining its high complement activity. When used in conjunction with Cytotoxicity Medium, this reagent provides a highly potent source of complement with minimal background toxicity.

Antibody Titration:

See **FIGURE 2**

Cell Source: spleen

Donor: C57BL/6

Cell Concentration: 1.1x10e6 cells/ml

Complement: Low-Tox®-M Rabbit Complement

Complement Concentration: 1:12

Procedure: Two stage cytotoxicity as described in Recommended Method for Determining Percent of Qa.m2 Positive Lymphocytes.

Strain Distribution:

Procedure: As above

Antibody Concentration: 1:40

Strains tested: C57BL/6, BALB/c, C3H/He,CBA, DBA/2,AKR

Cells killed by treatment: C57BL/6, DBA/2

Cells not killed by treatment: BALB/c, C3H/He, CBA, AKR

Tissue Distribution:

Procedure: as above

Antibody Concentration: 1:40

Strain: C57BL/6

Cell Source C.I.

Thymus: 10

Spleen: 43

Lymph Node: 64

Bone Marrow: 12

FLOW CYTOMETRY ANALYSIS:

See **FIGURE 4.**

DONOR: C57BL/6

CELL CONCENTRATION: 1 x 10e6 cells per test.

ANTIBODY CONCENTRATION: 1:500


CELL SOURCE: Splenic T Cells

Method

1. Prepare cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte®-M cell separation medium.
2. Wash 2 times.
3. Resuspend cells to 1×10^6 cells in approximately 50 μ l media A in a microcentrifuge tube (ie. 50 μ l of cells resuspended to 2×10^7 cells / ml). The contents of one tube represent one test.
4. To each tube add 50 μ l of a 1/250 dilution of this Ab (to make a final dilution of 1/500).
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Add 100 μ l of secondary antibody (Goat anti-mouse IgG (H+L) FITC conjugate) at a 1/500 dilution.
9. Incubate tubes at 4°C for 30-60 minutes.
10. It is recommended that the tubes are protected from light since most fluorochromes are light sensitive.
11. Wash 2 times at 4°C in media B.
12. Resuspend the cell pellet in 50 μ l ice cold media B.
13. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in phosphate buffered saline - (this stains dead cells by intercalating in DNA). N.B. Appropriate control samples should always be included in any labelling studies.

Media:

- A. Phosphate buffered saline (pH 7.2) + 0.5% normal serum of host species + sodium azide (100 μ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% bovine serum albumin + sodium azide (100 μ l of 2M sodium azide in 100 mls).

Product images:

LFL1

Figure 4

C.I. = Cytotoxic Index =
$$\frac{100 \times \% \text{cyt (antibody + complement)} - \% \text{cyt (complement alone)}}{100 \% - \% \text{cyt (complement alone)}}$$

Figure 1 - Cytotoxic Index

Strain	Qa.m2	Qa.m7	Qa.m8	Qa.m9
C57BL/6	+	+	+	+
A.BY	-	+	+	+
129	+	+	+	+
C3H	-	+	-	-
BIO.D2	+	+	-	+
BIO.M	-	-	-	-
BIO.BR	-	-	-	-
BIO.RIII(71NS)	-	-	-	-
BIO.S	+	+	-	+
A	+	-	-	-
CBA	-	-	-	-
C58	-	-	-	-
CE	-	-	-	-
NZB	+	+	+	-
DBA/1	+	+/-	-	-
DBA/2	+	+	-	-
SWR	+	+	-	-
BALB/cBy	-	-	-	-
SJL	+	+	+	-
RJII	-	-	-	ND
AKR	-	-	-	-
A.TL	+	-	-	ND

Figure 3: Strain Distribution of Qa.m2, Qa.m7, Qa.m8 and Qa.m9 specificities detected by monoclonal antibodies. - Protein Concentration: 45.8 ± 0.4 mg/ml




Figure 2