

Product datasheet for **CL072P**

MHC Class II I-Ap Mouse Monoclonal Antibody [Clone ID: 7-16.17]

Product data:

Product Type:	Primary Antibodies
Clone Name:	7-16.17
Applications:	CT, FC
Recommended Dilution:	Flow cytometry (protocol see below). Cytotoxic assay.
Reactivity:	Mouse
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	B10.p
Specificity:	CL072P is specific for Mouse-I-Ap,k,q,r,s,b. This antibody is a cytotoxic antibody which defines a public I-A antigen. Using recombinant strains, reactivity against the b haplotype has been localized to the Ab subregion. The antibody can be used to quantitate or eliminate I-A bearing cells for precipitating I-A antigen.
Formulation:	PBS and 0.02% sodium azide (NaN ₃). State: Purified State: Liquid Ig fraction
Concentration:	lot specific
Purification:	Protein G chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody at 2 - 8 °C for up to one month. For long term storage, aliquot and freeze unused portion at -20 °C in volumes appropriate for single usage. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.



[View online »](#)

Note: Protocol: **FLOW CYTOMETRY ANALYSIS:**

Method:

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population.
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in media A. Add 50 μ l of this suspension to each tube (each tube will then contain 1×10^6 cells, representing 1 test).
4. To each tube, add 0.2-1.0 μ g of antibody.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.
7. Wash 2 times at 4°C.
8. Add 100 μ l of secondary antibody (FITC Goat anti-mouse IgG) at 1/500 dilution.
9. Incubate the tubes at 4°C for 30-60 minutes. (It is recommended that the tubes are protected from light since most fluorochromes are light sensitive).
10. Wash 2 times at 4°C in media B.
11. Resuspend the cell pellet in 50 μ l ice cold media B.
12. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 μ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 μ l of 2M sodium azide in 100 mls).

Strain Distribution by Flow Cytometry Analysis:

Procedure: As above

Antibody Concentration: 0.5 μ g/ 10^6 cells

Strains Tested: see picture below.

For a more detailed strain distribution - see reference 1.