

## Product datasheet for **CL129F**

### MHC Class I (RT1Ac) Mouse Monoclonal Antibody [Clone ID: OX-27]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OX-27
Applications:	FC
Recommended Dilution:	Flow cytometry: use 0.5 µg of neat antibody to label 10e6 cells.
Reactivity:	Rat
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Phytohaemagglutinin Blasts.
Formulation:	PBS buffer with 0.02% sodium azide as preservative and 0.5 % EIA grade BSA as stabilizer. Label: FITC State: Liquid purified IgG fraction. Label: conjugated
Concentration:	lot specific
Purification:	Protein G Chromatography.
Conjugation:	FITC
Storage:	Store the antibody at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Database Link:	<a href="#">Q31255</a>
Background:	MHC Class I molecules play a central role in the immune system by presenting peptides derived from the endoplasmic reticulum lumen. MHC class I antigens are heterodimers consisting of one alpha chain (44kDa) with beta 2 microglobulin (11.5 kDa). The antigen is expressed by all somatic cells at varying levels. MHC Class I molecules are expressed on most nucleated cells where they present endogenously synthesized antigenic peptides to CD8+ T lymphocytes, which are usually cytotoxic T cells. Fibroblasts or neurons however only show a low level of antigen.
Synonyms:	RT1-A3



[View online »](#)

**Note:** Recognizes a polymorphic determinant of the MHC class I antigen in the rat. This Antibody can be used for labelling cells of donor or host origin in bone marrow chimeras. (1,2)

Protocol: **FLOW CYTOMETRY ANALYSIS:**

**Method:**

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Rat cell separation medium
2. Wash 2 times.
3. Resuspend the cells to a concentration of  $2 \times 10^7$  cells/ml in media A. Add 50  $\mu$ l of this suspension to each tube (each tube will then contain  $1 \times 10^6$  cells, representing 1 test).
4. To each tube, add 0.5-0.2  $\mu$ g antibody per  $10^6$  cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C. (protect tubes from light)
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50  $\mu$ l ice cold media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15  $\mu$ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

**Media:**

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100  $\mu$ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100  $\mu$ l of 2M sodium azide in 100 mls).

**Results-Tissue Distribution:**

Rat Strain: Brown Norway  
Cell Concentration:  $1 \times 10^6$  cells per test.  
Antibody Concentration Used: 0.2  $\mu$ g/ $10^6$  cells  
Isotypic Control: FITC Mouse IgG2a.

**Cell Source-Percentage of cells stained above control:**

Thymus: 73.9%  
Spleen: 99.0%  
Lymph Node: 100%