

# Product datasheet for DA3511X

### **KITLG / SCF Human Protein**

#### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


Product Type:	Recombinant Proteins
Description:	KITLG / SCF human recombinant protein, 5 μg
Species:	Human
Expression Host:	Insect
Predicted MW:	18.4 kDa
Purity:	>95% 95% by SDS-PAGE and visualised by silver stain
Buffer:	Presentation State: Purified State: Lyophilized purified protein Buffer System: PBS, pH 7.4, without stabilizer
Bioactivity:	Biological: Measured in a cell proliferation assay using TF 1 human erythroleukemic cells [Kitamura T et al, J Cell Physiol, 1989]. The ED50 for this effect is typically 1-5 ng/ml.
Endotoxin:	< 0.1 ng per µg of SCF
Reconstitution Method:	Restore in water to a concentration of 0.1 mg/ml. This solution can be diluted in water or other buffer solutions or stored at -20°C.
Preparation:	Lyophilized purified protein
Protein Description:	Recombinant Human Stem Cell Factor (SCF) His-tag. Soluble Stem Cell Factor (SCF), a 18.4kDa protein consisting of 165 amino acid residues (Glu26-Ala190) and fused to a C-terminal His-tag (6x His).
Storage:	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
RefSeq:	<u>NP 000890</u>
Locus ID:	4254
UniProt ID:	P21583, A0A024RBC0
Cytogenetics:	12q21.32
Synonyms:	DCUA; DFNA69; FPH2; FPHH; Kitl; KL-1; MGF; SCF; SF; SHEP7; SLF



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	KITLG / SCF Human Protein – DA3511X
Summary:	This gene encodes the ligand of the tyrosine-kinase receptor encoded by the KIT locus. This ligand is a pleiotropic factor that acts in utero in germ cell and neural cell development, and hematopoiesis, all believed to reflect a role in cell migration. In adults, it functions pleiotropically, while mostly noted for its continued requirement in hematopoiesis. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]
Protein Families:	Druggable Genome, Transmembrane
Protein Pathway	<b>s:</b> Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Melanogenesis, Pathways in cancer

## Product images:



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US