

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for DDX0010A546-100

CLEC10A Mouse Monoclonal Antibody [Clone ID: 125A10.03]

Product data:

Product Type:	Primary Antibodies
Clone Name:	125A10.03
Applications:	IF
Recommend Dilution:	IF
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	human ASGPR lg fusion protein
Specificity:	human ASGPR/MGL
Formulation:	Label: Alexa Fluor 546 State: 100 μg in 200 μl / 50 μg in 100 μl PBS 50% glycerol
Concentration:	0.5 mg/ml
Conjugation:	Alexa Fluor 546
Gene Name:	C-type lectin domain family 10 member A
Database Link:	Entrez Gene 10462 Human



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


CLEC10A Mouse Monoclonal Antibody [Clone ID: 125A10.03] – DDX0010A546-100

Background:ASGPR/MGL (asialoglycoprotein receptor/ macrophage galactose lectin receptor) / CD301 is a
Ca2+ dependent type II transmembrane lectin. In search for genes specifically expressed by
dendritic cells (DC), we have cloned several cDNAs encoding different forms of
asialoglycoprotein receptor (ASGPR), that were expressed in eukaryotic cells for use as
immunogens. Immunoprecipitation from DCs using anti-DCASGPR mAb yields a major 40-kDa
protein. mAb 125A10.03 recognized the ASGPR isoform expressed by DC but not the isoform
expressed by monocytes. This antibody stained CD14+derived DCs obtained from CD34+
progenitors, but not the CD1a+derived subset. Accordingly, both monocyte-derived DCs and
tonsillar interstitial-type DCs express DC-ASGPR protein, while Langerhans-type cells do not.
DC-ASGPR is a feature of immaturity, as its expression is lost upon CD40L activation. In
agreement with the presence of tyrosine-based and dileucine motifs in the intracytoplasmic
domain, mAb against DC-ASGPR is rapidly internalized by DCs at 37°C. Finally, intracellular
DC-ASGPR is localized in early endosomes, suggesting that the receptor recycles to the cell
surface following internalization of ligand. (*Valladeau J et al, 2001, J. Immunol., 167: 5767*).

Synonyms:

CLECSF13, CLECSF14, HML

Product images:



FACS analysisof monocyte-derivedDCs (GM+IL4) stained with 125A10.03+ goat anti-mouse FITC

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US