

## Product datasheet for **DDX0010A647-100**

### CLEC10A Mouse Monoclonal Antibody [Clone ID: 125A10.03]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	125A10.03
Applications:	FC
Recommend Dilution:	<b>Surface and intracyto Flow Cytometry</b>
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	human ASGPR Ig fusion protein
Specificity:	human ASGPR/MGL
Formulation:	Label: Alexa Fluor 647 State: 100 µg in 200 µl / 50 µg in 100 µl PBS 50% glycerol
Concentration:	0.5 mg/ml
Conjugation:	Alexa Fluor 647
Gene Name:	C-type lectin domain family 10 member A
Database Link:	<a href="#">Entrez Gene 10462 Human</a>




[View online »](#)

**Background:**

ASGPR/MGL (asialoglycoprotein receptor/ macrophage galactose lectin receptor) / CD301 is a  $\text{Ca}^{2+}$  dependent type II transmembrane lectin. In search for genes specifically expressed by dendritic cells (DC), we have cloned several cDNAs encoding different forms of asialoglycoprotein receptor (ASGPR), that were expressed in eukaryotic cells for use as immunogens. Immunoprecipitation from DCs using anti-DCASGPR mAb yields a major 40-kDa protein. mAb 125A10.03 recognized the ASGPR isoform expressed by DC but not the isoform expressed by monocytes. This antibody stained  $\text{CD14}^+$  derived DCs obtained from  $\text{CD34}^+$  progenitors, but not the  $\text{CD1a}^+$  derived subset. Accordingly, both monocyte-derived DCs and tonsillar interstitial-type DCs express DC-ASGPR protein, while Langerhans-type cells do not. DC-ASGPR is a feature of immaturity, as its expression is lost upon CD40L activation. In agreement with the presence of tyrosine-based and dileucine motifs in the intracytoplasmic domain, mAb against DC-ASGPR is rapidly internalized by DCs at  $37^\circ\text{C}$ . Finally, intracellular DC-ASGPR is localized in early endosomes, suggesting that the receptor recycles to the cell surface following internalization of ligand. (Valladeau *J et al*, 2001, *J. Immunol.*, 167: 5767).

**Synonyms:**

CLECSF13, CLECSF14, HML

**Product images:**

FACS analysis of monocyte-derived DCs (GM+IL4) stained with 125A10.03+ goat anti-mouse FITC