

## Product datasheet for DDX0011A488-100

## **CLEC10A Mouse Monoclonal Antibody [Clone ID: 121A5.01]**

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: 121A5.01
Applications: FC, IF

Recommend Dilution: Surface and intracyto Flow Cytometry, IF

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** human ASGPR Ig fusion protein

Specificity: human ASGPR/MGL
Formulation: Label: Alexa Fluor 488

State: 100 μg in 200 μl / 50μg in 100 μl PBS 50% glycerol

**Concentration:** 0.5 mg/ml

**Purification:** QMA Hyper D ion exchange chromatography

**Conjugation:** Alexa Fluor 488

**Gene Name:** C-type lectin domain family 10 member A

Database Link: Entrez Gene 10462 Human

OriGene Technologies, Inc.

Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com

EU: info-de@origene.com CN: techsupport@origene.cn

9620 Medical Center Drive, Ste 200




Background:

ASGPR/MGL (asialoglycoprotein receptor / macrophage galactose lectin receptor) / CD301 is a Ca2+ dependent type II transmembrane lectin. In search for genes specifically expressed by dendritic cells (DC), we have cloned several cDNAs encoding different forms of asialoglycoprotein receptor (ASGPR), that were expressed in eukaryotic cells for use as immunogens. Immunoprecipitation from DCs using anti-DCASGPR mAb yields a major 40-kDa protein. mAb 121A5.01 recognized the ASGPR isoform expressed by DC but not the isoform expressed by monocytes. This antibody stained CD14+derived DCs obtained from CD34+ progenitors, but not the CD1a+derived subset. Accordingly, both monocyte-derived DCs and tonsillar interstitial-type DCs express DC-ASGPR protein, while Langerhans-type cells do not. DC-ASGPR is a feature of immaturity, as its expression is lost upon CD40L activation. In agreement with the presence of tyrosine-based and dileucine motifs in the intracytoplasmic domain, mAb against DC-ASGPR is rapidly internalized by DCs at 37°C. Finally, intracellular DC-ASGPR is localized in early endosomes, suggesting that the receptor recycles to the cell surface following internalization of ligand. (*Valladeau J et al, 2001, J. Immunol., 167: 5767*).

Synonyms: CLECSF13, CLECSF14, HML

## **Product images:**



Monocyte-derived DC (GM+IL4) stained with 121A5.01+GAMFITC