

Product datasheet for **DDX0081A488-100**


CD1 (CD1A) Mouse Monoclonal Antibody [Clone ID: 201B5.08]

Product data:


Product Type:	Primary Antibodies
Clone Name:	201B5.08
Applications:	FC, IF
Recommend Dilution:	<u>DDX0081A488-50 / DDX0081A488-100 Alexa-fluor®488</u> : Surface flow cytometry, IF. <u>DDX0081A546-50 / DDX0081A546-100 Alexa- fluor®546 (on request)</u> : surface Flow Cytometry. <u>DDX0081A647-50 / DDX0081A647-100 Alexa- fluor®647</u> : Surface flow cytometry.
Reactivity:	Human, Porcine
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	in vitro derived human DCs (GMCSF + TNF α)
Specificity:	Human CD1a (epitope in extracellular domain). Species cross- reactivity: Pig.
Formulation:	Label: Alexa Fluor 488
Concentration:	0.5 mg/ml
Purification:	QMA hyper D ion exchange chromatography
Conjugation:	Alexa Fluor 488
Gene Name:	CD1a molecule
Database Link:	Entrez Gene 909 Human
Background:	CD1 family molecules have structural homologies with MHC class I molecules. They are monomeric proteins of 50 kDa, non-covalently associated with Beta-2-microglobulin. Human CD1a molecules are divided in 2 groups. Group 1 includes CD1a, CD1b, and CD1c, while group 2 consists of the CD1d molecule. CD1a is mainly expressed at the cell-surface and in recycling vesicles. CD1a occurs in Birbeck granules of Langerhans cells and plays a role in presentation of non-peptide antigens. CD1a is expressed in vitro on CD34 ⁺ -generated DCs and on monocyte-derived DCs. (<i>Burdin N. et al., 1999, Curr Opin Immunol, 11:326-31</i>).
Synonyms:	T6/Leu-6, hTa1



[View online »](#)

Product images:

IHC staining of human skin frozen section with clone 201B5 (DDX0081)



Surface FACS analysis of CD1a expression on CD34+-derived DCs (day 12 of culture)