

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for DDX0151A488-100

ICAM1 Mouse Monoclonal Antibody [Clone ID: 217E2.10]

Product data:

Product Type: Primary Antibodies

Clone Name: 217E2.10
Applications: FC, IF

Recommend Dilution: DDX0151P-50 / DDX0151P-100 Purified: Flow cytometry, IHC (cryosection, paraffin), IF.

<u>DDX0151A488-50 / DDX0151A488-100</u> Alexa-fluor® 488: Flow cytometry, IF (On request). <u>DDX0151A546-50 / DDX0151A546-100</u> Alexa- fluor® 546: Flow cytometry, IF (On request). <u>DDX0151A647-50 / DDX0151A647-100</u> Alexa- fluor® 647: Flow cytometry, IF (On request).

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: in vitro-derived DCs (GM-CSF+TNF)

Formulation:

Purified: 100 μ g in 200 μ l / 50 μ g in 100 μ l Tris-NaCl pH 8. **Coupled:** 100 μ g in 200 μ l / 50 μ g in 100 μ l PBS 50% glycerol

Label: Alexa Fluor 488

Concentration: 0.5 mg/ml

Conjugation: Alexa Fluor 488

Gene Name: intercellular adhesion molecule 1

Database Link: Entrez Gene 3383 Human

Background: 217E2 monoclonal antibody was obtained following mouse immunization with in vitro-

derived human dendritic cells. The target antigen was found to be CD54/ICAM-1, a cell surface glycoprotein which is typically expressed on endothelial cells and immune cells. CD54/ICAM-1, member of the Ig superfamily, binds to integrins of type CD11a / CD18, or CD11b / CD18 and is also a receptor for rhinovirus. CD54/ICAM-1 is a type of intracellular adhesion molecule, playing a role in transmigration of endothelial cells into tissues through

binding to LFA-1.


(Carlson M.et al, 1988, Nucleic Acids Res. 16 (9): 4188; Yang L. et al, 2005, Blood 106 (2): 584).

Synonyms: ICAM-1





Product images:



IHC staining of human lung frozen section with clone 217E2 (DX0151)