

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for DDX0195A488-100

## LAMP3 Mouse Monoclonal Antibody [Clone ID: 208B5]

## **Product data:**


Product Type:	Primary Antibodies
Clone Name:	208B5
Applications:	FC
Recommend Dilution:	<ul> <li>DDX0195P-50 / DDX0195P-100 Purified: FACS intracellular, ImmunoHistoChemistry frozen sections.</li> <li>DDX0195A488-50 / DDX0195A488-100 Alexa-fluor®488: Intra Flow cytometry for mouse and human.</li> <li>DDX0195A546-50 / DDX0195A546-100 Alexa-fluor®546: FACS intracellular, ImmunoHistoChemistry frozen sections.</li> <li>DDX0195A647-50 / DDX0195A647-100 Alexa-fluor®647: Intra Flow cytometry for mouse and human.</li> <li>DDX0195B-50 / DDX0195B-100 Biotin: FACS intracellular, ImmunoHistoChemistry frozen sections.</li> </ul>
	<b>Usage recommendation:</b> *This monoclonal antibody may be used between 5-10 $\mu$ g/ml.
Reactivity:	Human, Mouse
Host:	Mouse
lsotype:	lgG3
Clonality:	Monoclonal
Immunogen:	Recombinant Murine DC-LAMP
Specificity:	Murine DC-LAMP (epitope in intracytoplasmic domain). <b>Species cross-reactivity:</b> Human.
Formulation:	<u>Purified:</u> 100 μg in 200 μl / 50 μg in 100 μl Tris-NaCl pH 8. <u>Coupled</u> : 100 μg in 200 μl / 50 μg in 100 μl PBS 50% glycerol. Label: Alexa Fluor 488
Concentration:	0.5 mg/ml
Purification:	QMA Hyper D Ion exchange chromatography
Conjugation:	Alexa Fluor 488
Gene Name:	lysosomal associated membrane protein 3



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	LAMP3 Mouse Monoclonal Antibody [Clone ID: 208B5] – DDX0195A488-100
Database Link:	Entrez Gene 27074 Human
Background:	The dendritic cell lysosomal-associated membrane protein (DC-LAMP)/CD208 is a type I integral transmembrane glycoprotein mostly homologous to CD68, of about 45 kDa in the mouse and 90 kDa in human (due to extensive glycosylation), with a bipartite C-terminal structure divided by a serine/proline rich region, a transmembrane domain and a conserved tyrosine-based lysosomal targeting motif in its cytoplasmic tail. DC-LAMP is specifically expressed by human DCs upon activation. However, its mouse counterpart has not be detected in mature DCs. DC-LAMP is constitutively expressed by mouse. Confocal and immunoelectron microscopy has shown that mouse DC-LAMP protein co-localizes with lbm180, a specific marker for the limiting membrane of lamellar bodies that contains lung surfactant protein B ( <i>Salaun B et al, 2004; Eur. J. Immunol., 33: 2619-29</i> ).
Synonyms:	DCLAMP, DC-LAMP, TSC403, Dendritic Cell Marker

## Product images:



Intracellular flow cytometry with anti-DC-LAMP at  $5\mu$ g/ml on human monocyte-derived DC GM+ IL4 (24h in LPS)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US