

## Product datasheet for **DDX0203A546-100**

### DC SIGN (CD209) Mouse Monoclonal Antibody [Clone ID: 103G2.07]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	103G2.07
Applications:	IF
Recommend Dilution:	<b><u>DDX0203P-50 / DDX0203P-100 Purified:</u></b> FACS surface, ImmunoHistoChemistry frozen sections, ImmunoHistoChemistry paraffin sections. <b><u>DDX0203A488-50 / DDX0203A488-100</u></b> Alexa-fluor®488: Surface Flow cytometry, Immunofluorescence. <b><u>DDX0203A546-50 / DDX0203A546-100</u></b> Alexa- fluor®546: Immunofluorescence. <b><u>DDX0203A647-50 / DDX0203A647-100</u></b> Alexa- fluor®647: Surface Flow cytometry <b><u>DDX0203B-50 / DDX0203B-100</u></b> Biotin (on request): FACS surface, ImmunoHistoChemistry frozen sections, ImmunoHistoChemistry paraffin sections. <b>Usage recommendation:</b> *This monoclonal antibody may be used between 5-20 µg/ml.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	HeLa cells stably transfected-with human DC-SIGN.
Specificity:	Human DC-SIGN. <b>Species cross- reactivity:</b> Weak expression on NIH3T3-transfected LSIGN.
Formulation:	<b><u>Purified:</u></b> 100 µg in 200µl / 50 µg in 100 µl Tris-NaCl pH 8. <b><u>Coupled:</u></b> 100 µg in 200µl / 50 µg in 100 µl PBS 50% glycerol. Label: Alexa Fluor 546
Concentration:	0.5 mg/ml
Conjugation:	Alexa Fluor 546
Gene Name:	CD209 molecule
Database Link:	<a href="#">Entrez Gene 30835 Human</a>




[View online »](#)

**Background:**

DC-SIGN (“DC Specific, ICAM-3 Grabbing, Nonintegrin”)/CD209 and liver/lymph node-specific ICAM-3-grabbing nonintegrin (L-SIGN) (CD299/DC-SIGNR for DC-SIGN-related molecule; DC-SIGN2) are closely related genes that map to chromosome 19p13.3. Both genes encode a member of the C-type lectin family of type II transmembrane proteins. The two receptors are 77% identical at the amino acid level, have similar ligands. They are expressed in different tissues. DC-SIGN is expressed on dendritic cells and macrophages. L-SIGN is found in the endothelial cells of liver, lymph nodes, and placenta and is absent on DCs and macrophages. Both receptors have been shown to interact with ICAM-3 DC-SIGN is a high affinity receptor for HIV gp120 (Soilleux EJ. 2003, Clinical Science 104, 437-; Dakappagari N., et al. 2006, The J Immunol, 176, 426; Geijtenbeeck T.B., et al. 2000, Cell, 100, 57 ; .Bashirova A. et al., 2001, J.Exp. Med., 193, 671).

**Synonyms:**

DCSIGN1, DCSIGN, DC-SIGN1, CLEC4L, Dendritic Cell Marker

**Product images:**

Formalin-fixed paraffin-embedded human tonsil stained with clone 103G2 (DX0203). DC-SIGN is visualized by brown labeling with DAB