

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for DDX0363A488-100

Langerin (CD207) Mouse Monoclonal Antibody [Clone ID: DCGM4/122D5.03]

Product data:

Product Type: Primary Antibodies
Clone Name: DCGM4/122D5.03

Applications: FC, IF

Recommended Dilution: <u>DDX0363P-50 DDX0363P-100</u> **Purified:** Internalization, Flow cytometry, IHC.

DDX0363A488-50 DDX0363A488-100 Alexa-fluor®488: Flow cytometry, IF.

DDX0363A546-50 DDX0363A546-100 Alexa-fluor®546: IF.

DDX0363A647-50 DDX0363A647-100 Alexa- fluor®647: Flow cytometry.

DDX0363B-50 DDX0363B-100 Biotin: Western blot.

Usage recommendation:

*This monoclonal antibody may be used between 5-20 µg/ml.

*Optimal dilution should be determined y each laboratory for each application.

*Coupled antibody: to maintain RT before using.

Reactivity: Canine, Human

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: In vitro-derived human DCs from CD34+ cells in GM-CSF+TNFα

Specificity: Human Langerin (extracellular epitope in Carbohydrate Recognition domain).

Species cross-reactivity: Dog.

Formulation: Purified: 100 μg in 200μl / 50 μg in 100 μl Tris-NaCl pH 8.

Coupled: 100 µg in 200µl / 50 µg in 100 µl PBS 50% glycerol.

Label: Alexa Fluor 488

Concentration: lot specific

Purification: QMA Hyper D ion exchange chromatography

Conjugation: Alexa Fluor 488

Storage: -20°C. KEEP CONTENTS STERILE: no preservative.

<u>Purified antibodies</u>: avoid repeated freeze/thaw cycles. <u>Coupled antibodies</u>: glycerol protects from freezing.





Gene Name: CD207 molecule

Database Link: Entrez Gene 50489 Human


Background: Langerin/CD207 is a transmembrane C-type lectin receptor (CLR) of epidermal and mucosal

Langerhans cells (LCs) that induces Birbeck's granule formation. Langerin features a single carbohydrate recognition domain (CRD) with mannose-type specificity in its extracellular portion. Langerin is unique among the CLRs in that it contains an intracellular domain with a proline-rich motif. Langerin expression has not been reported outside the DC system. (Valladeau J et al, 1999, Eur.J.Immunol., 29:2695-2704; Valladeau J et al, 2000 Immunity, 12:71-81; Kashihara M et al, 1986, J.Invest.Derm., 87:602-607; Ito T et al, 1999, J.Immunol., 163:1409-1419 ;Saeland S & Valladeau J, CD207 (Langerin) Workshop reports 2002, Leukocyte-Typing VII,

White Cell Diff Antigens, D. Mason et al, Eds, Oxford University Press:306-307)

CLEC4K, Dendritic Cell Marker Synonyms:

Product images:



IHC staining of human skin frozen section with clone 122D5 (DX0363)






In vitro




Human skin

DCGM4(blue)



Western-blot DCGM4 1: non-reduced 2: reduced



sIgD (red) DCGM4(blue)

Valladeau J et al, 1999, Eur.J.Immunol