

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for DDX0363B-100

### Langerin (CD207) Mouse Monoclonal Antibody [Clone ID: DCGM4/122D5.03]

# Product data:

Product Type:	Primary Antibodies
Clone Name:	DCGM4/122D5.03
Applications:	FC, IHC, WB
Recommended Dilution:	DDX0363P-50 DDX0363P-100 Purified: Internalization, Flow cytometry, IHC. DDX0363A488-50 DDX0363A488-100 Alexa-fluor®488: Flow cytometry, IF. DDX0363A546-50 DDX0363A546-100 Alexa-fluor®546: IF. DDX0363A647-50 DDX0363A647-100 Alexa- fluor®647: Flow cytometry. DDX0363B-50 DDX0363B-100 Biotin: Western blot.
	<b>Usage recommendation:</b> *This monoclonal antibody may be used between 5-20 µg/ml. *Optimal dilution should be determined y each laboratory for each application. *Coupled antibody: to maintain RT before using.
Reactivity:	Canine, Human
Host:	Mouse
lsotype:	IgG1
Clonality:	Monoclonal
Immunogen:	In vitro-derived human DCs from CD34+ cells in GM-CSF+TNF $lpha$
Specificity:	Human Langerin (extracellular epitope in Carbohydrate Recognition domain). <b>Species cross-reactivity:</b> Dog.
Formulation:	<u>Purified:</u> 100 μg in 200μl / 50 μg in 100 μl Tris-NaCl pH 8. <u>Coupled:</u> 100 μg in 200μl / 50 μg in 100 μl PBS 50% glycerol. Label: Biotin
Concentration:	lot specific
Purification:	QMA Hyper D ion exchange chromatography
Conjugation:	Biotin
Storage:	-20°C. KEEP CONTENTS STERILE: no preservative. <u>Purified antibodies</u> : avoid repeated freeze/thaw cycles. <u>Coupled antibodies</u> : glycerol protects from freezing.




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2021 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Langerin (CD207) Mouse Monoclonal Antibody [Clone ID: DCGM4/122D5.03] – DDX0363B-100
Gene Name:	CD207 molecule
Database Link:	Entrez Gene 50489 Human
Background:	Langerin/CD207 is a transmembrane C-type lectin receptor (CLR) of epidermal and mucosal Langerhans cells (LCs) that induces Birbeck's granule formation. Langerin features a single carbohydrate recognition domain (CRD) with mannose-type specificity in its extracellular portion. Langerin is unique among the CLRs in that it contains an intracellular domain with a proline-rich motif. Langerin expression has not been reported outside the DC system. ( <i>Valladeau J et al, 1999, Eur.J.Immunol., 29:2695-2704; Valladeau J et al, 2000 Immunity, 12 : 71-</i> <i>81; Kashihara M et al, 1986, J.Invest.Derm., 87 :602-607 ; Ito T et al, 1999, J.Immunol., 163 :1409-</i> <i>1419 ;Saeland S &amp; Valladeau J, CD207 (Langerin) Workshop reports 2002, Leukocyte-Typing VII,</i> <i>White Cell Diff Antigens, D. Mason et al, Eds, Oxford University Press:306-307)</i>
-	

#### Synonyms:


#### CLEC4K, Dendritic Cell Marker

## **Product images:**




IHC staining of human skin frozen section with clone 122D5 (DX0363)





DCGM4(blue)



Western-blot DCGM4 1: non-reduced 2: reduced Valladeau J et al, 1999, Eur.J.Immunol



sIgD(red)DCGM4(blue)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2021 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US