

Product datasheet for **DDX0368A647-100**

Langerin (CD207) Mouse Monoclonal Antibody [Clone ID: 306G9.01/HD24]

Product data:

Product Type: Primary Antibodies
Clone Name: 306G9.01/HD24
Applications: FC
Recommend Dilution: **Flow cytometry, IF, IHC.**

Usage recommendation:

*This monoclonal antibody may be used between 5-20 µg/ml
*Optimal dilution should be determined by each laboratory for each application.
*Coupled antibody: to maintain RT before using.

Reactivity: Human, Mouse, Porcine, Rat
Host: Mouse
Isotype: IgG1
Clonality: Monoclonal
Immunogen: Mouse langerin-transfected COP5 cells.
Specificity: Mouse and Human Langerin (intracellular epitope).
Species cross-reactivity: Human, Rat, Swine.

Formulation: **Purified:** 100 µg in 200µl / 50 µg in 100 µl Tris-NaCl pH 8.
Coupled: 100 µg in 200µl / 50 µg in 100 µl PBS 50% glycerol.
Label: Alexa Fluor 647

Concentration: 0.5 mg/ml
Purification: QMA Hyper D ion exchange chromatography
Conjugation: Alexa Fluor 647
Gene Name: CD207 molecule
Database Link: [Entrez Gene 50489 Human](#)



[View online »](#)


Background:

Langerin/CD207 is a transmembrane C-type lectin receptor (CLR) of epidermal and mucosal Langerhans cells (LCs) that induces Birbeck granule formation. Langerin features a single carbohydrate recognition domain (CRD) with mannose-type specificity in its extracellular portion. Langerin is unique among the CLRs in that it contains an intracellular domain with a proline-rich motif. Langerin expression has not been reported outside the DC system. *In mouse, clone 306G9 stains lymph node dendritic cells (Henri et al, 2001, J. Immunol, 167: 741-748) (Valladeau J. et al, 2002 J Immunol, 168: 782).*


Synonyms:

CLEC4K, Dendritic Cell Marker


Product images:




IHC staining of human skin frozen section with clone 306G9 (DX0368)




IHC staining of **human** skin frozen section with 306G9 antibody




IHC staining of **human** skin paraffin section with 306G9 antibody



Facs analysis of **mouse** Langerin-transfected COP5 cells stained with 306G9 antibody



IHC staining of Langerhans cells in **porcine** skin tissue section



Facs double-staining of **porcine** skin-derived DCs