

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for DDX0462A546-100

SEMA6A Mouse Monoclonal Antibody [Clone ID: 104B3]

Product data:

Product Type: Primary Antibodies

Clone Name: 104B3

Applications: IF

Recommend Dilution: <u>DDX0462P-50 / DDX0462P-100</u> Purified: IHC.

DDX0462A488-50 / DDX0462A488-100 Alexa-fluor®488: IF, Flow cytometry

DDX0462A546-50 / DDX0462A546-100 Alexa-fluor®546: IF

<u>DDX0462A647-50 / DDX0462A647-100</u> Alexa- fluor®647: (on request) IHC, Flow cytometry.

Usage recommendation:

*This monoclonal antibody may be used between 5-20 µg/ml.

*Optimal dilution should be determined by each laboratory for each application.

*Coupled antibody: to maintain RT before using.

Reactivity: Human

Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: human recombinant SEMA-6A

Specificity: Human SEMA-6A (epitope in extracellular domain).

Formulation: Purified: 100 µg in 200µl / 50 µg in 100 µl Tris-NaCl pH 8.

Coupled: 100 μg in 200μl / 50 μg in 100 μl PBS 50% glycerol.

Label: Alexa Fluor 546

Concentration: 0.5 mg/ml

Purification: QMA Hyper D ion exchange chromatography

Conjugation: Alexa Fluor 546
Gene Name: semaphorin 6A

Database Link: Entrez Gene 57556 Human






Background:

Semaphorins represent a large family composed of 19 secreted, GPI-linked or transmembrane members divided into 5 classes (3 to 7) and characterized by the presence of a 500 amino acid Sema domain in the Nterminus. Initially described for their role in axon guidance and apoptosis during neuronal development, some semaphorins like the class 4 regulate immune functions and are involved in organogenesis and angiogenesis. The transmembrane class 6 semaphorins include 4 members (SEMA-6A to 6D). SEMA-6A features an intracellular binding site for regulatory EvI protein. In the immune system, SEMA6A protein is restricted in vitro and in vivo to an activation status of Langerhans cells. (*Gautier G et al.*, 2006 Am J Pathol., 168: 453-65).


Synonyms:

Semaphorin-6A, Semaphorin VIA, Sema VIA, Semaphorin-6A-1, SEMA6A-1, SEMAQ, KIAA1368

Product images:



Gautier et al, 2006



IHC staining of Langerhans histiocytosis frozen section with clone 104B3 (DDX0462)