

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for DDX0710P-100

IL1 alpha (IL1A) Human Monoclonal Antibody [Clone ID: HuMab X3]

Product data:

Product Type: Primary Antibodies

Clone Name: HuMab X3

Applications: NEUT

Recommend Dilution: DDX0710P-50 / DDX0710P-100 Purified: Inhibition, Neutralization (Binding to IL1α and not to

IL1 or IL1Ra, Neutralizing biological effects of natural and recombinant IL1α, Inhibition of

IL1α binding to IL1 receptors.

<u>DDX0710A488-50 / DDX0710A488-100</u> Alexa-fluor®488 (on request). <u>DDX0710A546-50 / DDX0710A546-100</u> Alexa-fluor®546 (on request). <u>DDX0710A647-50 / DDX0710A647-100</u> Alexa-fluor®647 (on request).

<u>DDX0710B-50 / DDX0710B-100</u> Biotin (on request).

Usage recommendation:

*This monoclonal antibody may be used between 5-25 µg/ml.

*Optimal dilution should be determined by each laboratory for each application.

*Coupled antibody: to maintain RT before using.

Reactivity: Human
Host: Human
Isotype: IgG4

Clonality: Monoclonal

 $\begin{tabular}{ll} \textbf{Immunogen:} & CD40 \ activated \ and \ EBV \ immortalized \ human \ B \ lymphocytes. \\ \textbf{Specificity:} & Human \ IL1\alpha \ (binds \ to \ Human \ IL1\alpha \ but \ not \ to \ IL1\beta \ and \ IL1Ra). \\ \end{tabular}$

Formulation: Purified: 100 µg in 200µl / 50 µg in 100 µl Tris-NaCl pH 8.

Coupled: 100 μg in 200μl / 50 μg in 100 μl Tris PBS 50% glycerol.

State: Purified

Concentration: 0.5 mg/ml

Purification: QMA Hyper D ion exchange chromatography

Gene Name: interleukin 1 alpha

Database Link: Entrez Gene 3552 Human





Background:


Interleukin-1 (IL-1) defines two polypeptides, IL-1 α and IL-1 β , that possess a wide spectrum of biological effects. Two natural antagonists of IL-1 action have been characterized: the IL-1 receptor antagonist (IL-1Ra) and a soluble form of the type II IL-1 receptor. Neutralizing autoantibodies to IL-1 α have also been detected in sera of healthy individuals and patients with autoimmune or inflammatory diseases. By combining CD40-activation and Epstein-Barr virusimmortalization of B lymphocytes from individuals with circulating anti-IL-1 α , a natural anti-IL- 1 α monoclonal autoantibody was developed that bound specifically to human IL-1 α , but not to IL-1 β and IL-1Ra, with a high affinity (Kd = 1.2 × 10⁻¹⁰ M).

HuMAb X3 inhibited IL-1 α binding to IL-1 receptors and neutralized biological activities of both recombinant and natural forms of IL-1 α . This high affinity neutralizing anti-IL-1 α human autoantibody can function as a third type of IL-1 antagonist. (Garrone P et al, Mol Immunol 1996, 33 (7-8), 649-58; Guillot-Chene P, Lebecque S, Rigal D, Annales Pharmaceutiques Françaises, 06/2009, 67 (3) 182-6).

Synonyms:


IL-1 alpha, IL1F1, IL1A, Hematopoietin-1

Product images:




Inhibition of human IL-1 α -induced IL-6 production by human synoviocytes by HuMab X3





Inhibition of human IL-1 α -induced IL-6 production by human synoviocytes by HuMab X3 in the presence of either human IL-1 β or IL-1 α



"Cross-Reactivity of the HuMab X3" using excess human IL-1 β , IL-1Ra or IL-1 α to protect against immunoprecipitation byHuMAb X3.