

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for DDX0910P-100**

alpha Synuclein (SNCA) Mouse Monoclonal Antibody [Clone ID: AS11]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: AS11

Applications: IF, IHC, WB

Recommended Dilution: <u>DDX0910P-50 / DDX0910P-100</u> Purified: ImmunoFluorescence, ImmunoHistoChemistry

frozen sections, Western Blot.

<u>DDX0910A488-50 / DDX0910A488-100</u> Alexa-fluor®488: ImmunoFluorescence. <u>DDX0910A546-50 / DDX0910A546-100</u> Alexa-fluor®546: ImmunoFluorescence. <u>DDX0910A647-50 / DDX0910A647-100</u> Alexa-fluor®647: ImmunoFluorescence.

DDX0910HRP-50 / DDX0910HRP-100 Horseradish Peroxydase: ImmunoHistoChemistry

frozen sections, Western Blot.

**Usage recommendation:** 

\*This monoclonal antibody may be used between 5-20  $\mu$ g/ml.

\*Optimal dilution should be determined by each laboratory for each application.

\*Coupled antibody: to maintain RT before use.

Reactivity: Human
Host: Mouse

Isotype: lgG

Clonality: Monoclonal

**Immunogen:** rhu α-synuclein (SIGMA).

**Specificity:** N-terminal portion of Human  $\alpha$ -synuclein (aa1-60)

**Species cross-reactivity:** Negative for Murine  $\alpha$ -synuclein.

**Formulation:** Purified: 100 µg in 200µl / 50 µg in 100 µl Tris-NaCl pH 8.

**Coupled:** 100 μg in 200μl / 50 μg in 100 μl Tris PBS 50% glycerol.

State: Purified

**Concentration:** lot specific

**Purification:** QMA Hyper D ion exchange chromatography





-20°C. KEEP CONTENTS STERILE: no preservative. Storage:

> Purified antibodies: avoid repeated freeze/thaw cycles. **Coupled antibodies:** glycerol protects from freezing.

Gene Name: synuclein alpha

Database Link: Entrez Gene 6622 Human

Background: α-synuclein (snca) is a140 aas neuronal protein predominantly present in synaptic junctions,

specially studied for its relevance in understanding sporadic and familial forms of Parkinson's

disease and related neurodegenerative disorders. In an attempt to break the

anti-bovine immunodeficiency virus


(negative control)

immunotolerance since the amino-acid sequence is highly conserved between species (95.3% homology between human and mouse), we have immunized C57BL/6 inbred mice presenting a natural deletion for snca with recombinant human  $\alpha$ -synuclein and obtained a murine mAb (AS11) recognizing endogenous human α-synuclein. (Kahle et al., 2000, J Neurosci 2000; 20:6365-73; Specht and Schoepfer, BMC Neurosci 2001;2:11; Mougenot AL et al, | Neurosci Methods. 2010

Oct 15;192(2):268-76).

Synonyms: NACP, PARK1

## **Product images:**



anti-α-synuclein (Abcam)

IF staining of human neuroblastoma (SH-SY5Y) expressing endogenous α-synuclein

IHC staining of pathological human brainswith AS11