

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for DDX1401P-100

MYD88 Mouse Monoclonal Antibody [Clone ID: 603E10.06]

Product data:

Product Type: Primary Antibodies

Clone Name: 603E10.06

Applications: FC, IF

Recommended Dilution: <u>DDX1401P-50 / DDX1401P-100</u> Purified: FACS intracellular, ImmunoFluorescence.

DDX1401A488-50 / DDX1401A488-100 Alexa- fluor®A488: FACS intracellular,

Immuno Fluorescence.

DDX1401A546-50 / DDX1401A546-100 Alexa- fluor®546: FACS intracellular,

ImmunoFluorescence.

DDX1401A647-50 / DDX1401A647-100 Alexa- fluor®647: FACS intracellular,

ImmunoFluorescence.

Applications tested: Intracyto flow cytometry, IF, IHC paraffin.

Usage recommendation:

*This monoclonal antibody may be used between 5-20 µg/ml.

*Optimal dilution should be determined by each laboratory for each application.

*Coupled antibody: to maintain RT before using.

Reactivity: Canine, Human, Porcine

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: huMyd88 transfected 293T cells

Specificity: Human Myd88.

Both 603E10.05 and 603E10.06 monoclonal antibodies were validated both in Human and in

Dog. 603E10.05 was validated in Swine.

Species cross-reactivity: Negative for Murine Myd88.

Formulation: Purified: 100 µg in 200µl / 50 µg in 100 µl Tris-NaCl pH 8.

Coupled: 100 μg in 200μl / 50 μg in 100 μl Tris PBS 50% glycerol.

State: Purified

Concentration: lot specific

Purification: QMA Hyper D ion exchange chromatography





Storage: -20°C. KEEP CONTENTS STERILE: no preservative.

<u>Purified antibodies</u>: avoid repeated freeze/thaw cycles. <u>Coupled antibodies</u>: glycerol protects from freezing.

Gene Name: myeloid differentiation primary response 88

Database Link: Entrez Gene 4615 Human

Background: MyD88 (myeloid differentiation primary response gene 88) is a universal adapter protein of

296 aa used by all TLRs (except TLR3) and by IL1-R to activate the transcription factor NF-kB. Myd88 binds IRAK1, IRAK2 and TRAF6, leading to NF-kB activation, cytokine secretion and inflammatory response. Myd88 increases IL8 transcription, and is involved in IL18 signaling pathway. A monoclonal antibody was raised against huMyd88-transfected 293T cells. This antibody specifically recognizes endogenously expressed Myd88 and does not cross-react with the mouse Myd88. Both 603E10.05 and 603E10.06 monoclonal antibodies were

validated both in human and in dog. 603E10.05 was validated in swine.

Synonyms: Myeloid differentiation primary response protein MyD88

Product images:







