

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for DDX9022B-100

# CD49b (ITGA2) Human Monoclonal Antibody [Clone ID: B4F12]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	B4F12
Applications:	ELISA, FC
Recommended Dilution:	<ul> <li>IF, ELISA, preferential detection of activated platelet by flow cytometry, platelet aggregation studies.</li> <li>Usage recommendation:</li> <li>*This monoclonal antibody may be used between 5-25µg/ml.</li> <li>*Optimal dilution should be determined by each laboratory for each application.</li> <li>*Coupled antibody: to maintain RT before use.</li> </ul>
Reactivity:	Human
Host:	Human
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	CD40 activated and EBV immortalized human B lymphocytes
Specificity:	GPIa-IIa preferentially on acitvated platelets.
Formulation:	100 μg in 200 μl / 50 μg in 100 μl PBS 50% glycerol Label: Biotin
Concentration:	lot specific
Purification:	QMA Hyper D ion exchange chromatography
Conjugation:	Biotin
Storage:	-20°C. KEEP CONTENTS STERILE: no preservative. Purified antibodies: avoid repeated freeze/thaw cycles. Coupled antibodies: glycerol protects from freezing.
Gene Name:	integrin subunit alpha 2
Database Link:	Entrez Gene 3673 Human



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2021 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### **CD49b** (ITGA2) Human Monoclonal Antibody [Clone ID: B4F12] – DDX9022B-100

#### Background:


Neonatal Allolmmune Thrombocytopenia (NAIT) is the commonest cause of severe isolated thrombocytopenia in the foetus and newborn. In Caucasians, 97% of NAIT are caused by HPA-1a or HPA- 5b alloantibodies. B4F12 was isolated after combination of CD40 activation and Epstein-Barr virus transformation of B cells isolated from a woman with high titre of anti -HPA-5b alloantibodies. B4F12 human monoclonal antibody recognizes GPIa-IIa preferentially on activated human platelets (*Mérieux Y, Rigal D, Pin JJ, XVIIth Regional Congress, International Society of Blood Transfusion, Madrid 2007 ; Guillot-Chene P, Lebecque S, Rigal D, Annales Pharmaceutiques Françaises, 06/2009, 67 (3) 182-6*).

#### Synonyms:

Integrin alpha-2, ITGA-2, VLA-2 alpha, VLA2, GPIa, Collagen Receptor

% inhibition

### **Product images:**



IF staining of human platelets with **B4F12** 



ELISA detection of human platelets with **B4F12** 

mAb (ug/ml)

Inhibition of collagen-induced platelet aggregation by **F2H11** and **B4F12** 

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2021 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US