

## Product datasheet for DM055P

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cytokeratin 10 (KRT10) Mouse Monoclonal Antibody [Clone ID: DE-K10]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: DE-K10

**Applications:** FC, IF, IHC, WB

Recommended Dilution: Immunoblotting.

Flow Cytometry.

Immunocytochemistry.

Immunohistochemistry on Frozen Sections.

Immunohistochemistry on Paraffin-Embedded Sections.

Recommended Dilutions: 1/100–1/200 for Flow Cytometry, and for Immunohistochemistry with avidin-biotinylated horseradish peroxidase complex (ABC) as detection reagent, and

1/100–1/1000 for Immunoblotting applications.

**Reactivity:** Canine, Feline, Human

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Cytoskeletal preparation extracted from Human epidermis.

**Specificity:** DE-K10 reacts exclusively with cytokeratin 10 which is present in keratinizing stratified

epithelia and in differentiated areas of highly differentiated squamous cell carcinomas.

Formulation: PBS

State: Purified

State: Liquid purified IgG fraction Preservative: 0.09% Sodium Azide

**Concentration:** lot specific

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Gene Name:** keratin 10





## Cytokeratin 10 (KRT10) Mouse Monoclonal Antibody [Clone ID: DE-K10] - DM055P

Database Link: Entrez Gene 3858 Human

P13645

**Background:** Cytokeratins are a subfamily of intermediate filament proteins and are characterized by a

remarkable biochemical diversity, represented in human epithelial tissues by at least 20 different polypeptides. They range in molecular weight between 40 kDa and 68 kDa and isoelectric pH between 4.9 - 7.8. The individual human cytokeratins are numbered 1 to 20. The various epithelia in the human body usually express cytokeratins which are not only characteristic of the type of epithelium, but also related to the degree of maturation or

differentiation within an epithelium.

Cytokeratin subtype expression patterns are used to an increasing extent in the distinction of

different types of epithelial malignancies. The cytokeratin antibodies are not only of assistance in the differential diagnosis of tumors using immunohistochemistry on tissue

sections, but are also a useful tool in cytopathology and flow cytometric assays.

Synonyms: Cytokeratin-10, CK10, CK-10, Keratin-10, KRT10, KPP, K10