

Product datasheet for **DM1013**

GM CSF (CSF2) Mouse Monoclonal Antibody [Clone ID: 29]

Product data:

Product Type:	Primary Antibodies
Clone Name:	29
Applications:	ELISA, WB
Recommended Dilution:	ELISA. Western Blot.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Purified recombinant human GM-CSF Mouse IgG1.
Specificity:	This monoclonal anti-GM-CSF antibody reacts with natural and recombinant Human GM-CSF. Does not show any cross reaction with other human Cytokines or Growth Factors tested such as M-CSF, G-CSFR, IL-8, IL-16, IL1-beta, TGFbeta-1 and TNF-alpha.
Formulation:	0.01M PBS, pH 7.2 without preservatives. State: Purified State: Lyophilized purified IgG fraction.
Reconstitution Method:	Restore with double distilled water is recommended and to adjust the final concentration to 1.00 mg/ml
Concentration:	lot specific
Purification:	Affinity Chromatography on Protein G.
Conjugation:	Unconjugated
Storage:	Store the antibody at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	colony stimulating factor 2
Database Link:	Entrez Gene 1437 Human P04141



[View online »](#)

Background:	The cytokine GM-CSF stimulates the growth and differentiation of hematopoietic precursor cells from various lineages, including granulocytes, macrophages, eosinophils and erythrocytes. Used in myeloid reconstitution following bone marrow transplant, bone marrow transplant engraftment failure or delay, mobilization and following transplantation of autologous peripheral blood progenitor cells, and following induction chemotherapy in older adults with acute myelogenous leukemia.
Synonyms:	CSF2, GMCSF, Sargramostim, Molgramostin
Protein Families:	Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein
Protein Pathways:	Cytokine-cytokine receptor interaction, Fc epsilon RI signaling pathway, Hematopoietic cell lineage, Jak-STAT signaling pathway, Natural killer cell mediated cytotoxicity, T cell receptor signaling pathway