

Product datasheet for **DM1022**

IL8 (CXCL8) Mouse Monoclonal Antibody [Clone ID: I8-S2]

Product data:

Product Type:	Primary Antibodies
Clone Name:	I8-S2
Applications:	ELISA, FN, WB
Recommended Dilution:	ELISA: Reacts with human IL-8 (matched pair with clone I8-60, Cat.-No DM1023). Neutralizing: In chemotaxis assay, the antibody inhibited 97.8% % of the chemotactic activity of IL-8 on RB/293 cell using 5 µg/ml of MAb and 10 ng/ml of human IL-8 in assay. This MAb inhibited the chemotactic effect of human IL-8, had no inhibitory effect to MIP 1-beta and RANTES in neutrophil chemotaxis assay. It indicates the MAb is special neutralizing the IL-8 activity. The MAb has been shown the neutralizing activity on calcium ion changes in human granulocytes using FACS. Western Blot: Concentration of 0.02-0.1 µg/ml MAb will allow visualization of 100 ng/lane of human IL-8.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Purified recombinant Human IL-8.
Specificity:	This monoclonal antibody recognizes both recombinant and native Human Interleukin-8. This antibody does not cross react with human Monocyte Chemotactic Activating Factor (MCAF) or RANTES (Regulated on Activation, Normal T-cell Expressed, and Secreted).
Formulation:	0.01M PBS, pH 7.0 State: Azide Free State: Lyophilized purified Ig fraction Preservative: None
Reconstitution Method:	Restore with double distilled water to adjust the final concentration to 1.0 mg/ml.
Concentration:	1.0 mg/ml (after reconstitution)
Purification:	Affinity Chromatography on Protein G
Conjugation:	Unconjugated



[View online »](#)

Storage: Upon receipt, store (in aliquots) at -20°C.
Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.


Gene Name: C-X-C motif chemokine ligand 8

Database Link: [Entrez Gene 3576 Human P10145](#)

Background: Interleukin 8 (IL-8), formerly called monocyte-derived neutrophil chemotactic factor, belongs to the C-X-C chemokine family. IL-8 has 4 cysteine residues, as do other members of the chemokine family, and the first two cysteine residues are separated by glutamine. IL-8 consists of 72 amino acids with a molecular weight of 8,000 daltons. IL-8 exhibits chemotactic activity in vitro for T cells, basophils and neutrophils. IL-8 activates neutrophils to release lysosomal enzymes including myeloperoxidase, -mannosidase and -glucuronidase. IL-1 induces the production of IL-8 from fibroblasts, keratinocytes, endothelial cells, hepatoma cells, astrocytoma cells, glioblastoma cells, lung epithelial cells, synovial membrane cells, melanocytes, melanoma cells and gastric cancer cells. Lipopolysaccharides will stimulate IL-8 production in monocytes/macrophages, and expressed on endothelial cells in response to inflammation.

Synonyms: CXCL8, Protein 3-10C, Emoctakin, GCP1, MDNCF, MONAP, NAP1

Product images:



Western blot analysis of Human IL8 using anti-Human IL8 antibody