

Product datasheet for DM196-05

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Von Willebrand Factor (VWF) Mouse Monoclonal Antibody [Clone ID: F8/86]

Product data:

Product Type: Primary Antibodies

Clone Name: F8/86
Applications: IHC

Recommended Dilution: Immunohistochemistry on Paraffin Embedded Sections: Use Factor VIII R ag antibody at

1/15-1/30 dilution in an ABC method (30 minutes at RT).

Proteolytic enzyme digestion of formalin fixed paraffin embedded tissue sections is required

prior to immunostaining.

Recommended Positive Control: Tonsil.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Von Willebrand factor isolated from Human plasma.

Specificity: This antibody reacts with Von Willebrand factor in endothelial cells. It also reacts with

megakaryocytes in Human bone marrow.

Cellular Localization: Cytoplasmic.

Formulation: State: Purified

State: Liquid purified immunoglobulin fraction

Preservative: < 0.1% Sodium Azide

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: von Willebrand factor

Database Link: Entrez Gene 7450 Human


P04275

Synonyms: vWF, von Willebrand antigen 2, von Willebrand antigen II, F8VWF, Factor VIII Related Antigen





Product images:



Formalin fixed paraffin embedded human tonsil stained with Factor VIII antibody DM196