

Product datasheet for DM2045

CCL3 Mouse Monoclonal Antibody [Clone ID: 199]

Product data:

Isotype:

Product Type: Primary Antibodies

Clone Name: 199
Applications: ELISA
Recommended Dilution: ELISA.
Reactivity: Human
Host: Mouse

Clonality: Monoclonal

Immunogen: Purified recombinant Human MIP1 alpha.

IgG2a

Specificity: This antibody reacts with recombinant MIP1 alpha.

Does not show any cross reaction with other human cytokines or growth factors tested such

as M-CSF, GM-CSF, IL-7, IL-8, IL-16, IL1 beta, bFGF, TGF beta, EGF, TNF alpha and EPO.

Formulation: 0.01M PBS, pH 7.2 without preservatives.

State: Purified

State: Lyophilized purified IgG fraction.

Reconstitution Method: Restore with double distillated water to adjust the final concentration to 1.0 mg/ml.

Purification: Affinity Chromatography on Protein G.

Conjugation: Unconjugated

Storage: Store the antibody at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: C-C motif chemokine ligand 3

Database Link: Entrez Gene 6348 Human

P10147



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



CCL3 Mouse Monoclonal Antibody [Clone ID: 199] - DM2045

Background:

Macrophage Inflammatory Protein 1 alpha (MIP1 alpha) and MIP1 beta, two closely related but distinct proteins, were originally co-purified from medium conditioned by a LPS-stimulated murine macrophage cell line. Mature mouse MIP1 alpha shares approximately 77% and 70% amino acid identity with human MIP1 alpha and mouse MIP1 beta, respectively. MIP1 proteins are expressed primarily in T cells, B cells, and monocytes after antigen or mitogen stimulation. Has adhesive effects on lymphocytes. MIP1 alpha can inhibit the proliferation of hematopoietic stem cells in vitro as well as in vivo. A signal transducing receptor, designated the C-C chemokine receptor-1 (C-C CKR-1) with seven transmembrane domains that bind MIP1 alpha, MIP1 beta, MCP1 and RANTES with varying affinities, has been isolated.

Synonyms:

CCL-3, MIP-1-alpha, C-C motif chemokine 3, MIP1A, Small-inducible cytokine A3, SCYA3, LD78 alpha, G0S19-1 protein