

Product datasheet for DM2047

CCL3 Mouse Monoclonal Antibody [Clone ID: 400]

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Clone Name:	400
Applications:	ELISA, WB
Recommended Dilution:	ELISA. Western Blot.
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Purified recombinant Human MIP1 alpha.
Specificity:	This antibody reacts with recombinant MIP1 alpha. Does not show any cross reaction with other human cytokines or growth factors tested such as M-CSF, GM-CSF, IL-7, IL-8, IL-16, IL1 beta, bFGF, TGF beta, EGF, TNF alpha and EPO.
Formulation:	0.01M PBS, pH 7.2 without preservatives State: Purified State: Lyophilized purified IgG fraction.
Reconstitution Method:	Restore with double distillated water to adjust the final concentration to 1.0 mg/ml.
Purification:	Affinity Chromatography on Protein G.
Conjugation:	Unconjugated
Storage:	Store the antibody at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	C-C motif chemokine ligand 3
Database Link:	<u>Entrez Gene 6348 Human</u> <u>P10147</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	CCL3 Mouse Monoclonal Antibody [Clone ID: 400] – DM2047
Background:	Macrophage Inflammatory Protein 1 alpha (MIP1 alpha) and MIP1 beta, two closely related but distinct proteins, were originally co-purified from medium conditioned by a LPS- stimulated murine macrophage cell line. Mature mouse MIP1 alpha shares approximately 77% and 70% amino acid identity with human MIP1 alpha and mouse MIP1 beta, respectively. MIP1 proteins are expressed primarily in T cells, B cells, and monocytes after antigen or mitogen stimulation. Has adhesive effects on lymphocytes. MIP1 alpha can inhibit the proliferation of hematopoietic stem cells in vitro as well as in vivo. A signal transducing receptor, designated the C-C chemokine receptor-1 (C-C CKR-1) with seven transmembrane domains that bind MIP1 alpha, MIP1 beta, MCP1 and RANTES with varying affinities, has been isolated.
Synonyms:	CCL-3, MIP-1-alpha, C-C motif chemokine 3, MIP1A, Small-inducible cytokine A3, SCYA3, LD78 alpha, G0S19-1 protein
Protein Familie	s: Druggable Genome, Secreted Protein
Protein Pathwa	ys: Chemokine signaling pathway, Cytokine-cytokine receptor interaction, Toll-like receptor signaling pathway

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US