

Product datasheet for **DM3506**

VEGF Receptor 1 (FLT1) Mouse Monoclonal Antibody [Clone ID: FLTEWF]

Product data:

Product Type:	Primary Antibodies
Clone Name:	FLTEWF
Applications:	ELISA, IF, IP, WB
Recommended Dilution:	ELISA: 1-10 µg/ml. Western blotting: 1-10 µg/ml. Immunoprecipitation: 1-5 µg/ml lysate or reaction volume.
Reactivity:	Human, Mouse
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Recombinant Human soluble extracellular Flt-1 protein (D5) (Cat.-No DA3539)
Specificity:	The monoclonal antibody will recognize recombinant and naturally occurring form of Human VEGFR-1/Flt-1.
Formulation:	PBS pH 7.4 containing no preservative State: Purified State: Lyophilized purified IgG fraction
Reconstitution Method:	Restore with distilled sterile water to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity Chromatography on Protein G
Conjugation:	Unconjugated
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	fms related tyrosine kinase 1
Database Link:	Entrez Gene 2321 Human P17948





[View online »](#)

Background:

Endothelial cells express three different vascular endothelial growth factor (VEGF) receptors, belonging to the family of receptor tyrosine kinases (RTKs). They are named VEGFR-1 (Flt-1), VEGFR-2 (KDR/Flk-1), VEGFR-3 (Flt-4). Their expression is almost exclusively restricted to endothelial cells, but VEGFR-1 can also be found on monocytes, dendritic cells and on trophoblast cells. The flt-1 gene was first described in 1990. The receptor contains seven immunoglobulin-like extracellular domains, a single transmembrane region and an intracellular splitted tyrosine kinase domain. Compared to VEGFR-2 the Flt-1 receptor has a higher affinity for VEGF but a weaker signaling activity. VEGFR-1 thus leads not to proliferation of endothelial cells, but mediates signals for differentiation. Interestingly a naturally occurring soluble variant of VEGFR-1 (sVEGFR-1) was found in HUVE supernatants in 1996, which is generated by alternative splicing of the flt-1 mRNA.

Synonyms:

VEGFR1, FLT1, FLT, FRT, VEGF Receptor 1

Product images:


Standard ELISA assay with coating of 2.5 µg/ml Mouse sFlt-1-Fc using anti Human Flt-1 antibodies