

Product datasheet for **DP012-05**

Von Willebrand Factor (VWF) Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	IHC
Recommended Dilution:	Immunohistochemistry on formalin-fixed paraffin embedded tissues: This antibody may be diluted to a titer of 1:50-1:100 in an ABC method (30 minutes at room temperature). Although not required, proteolytic treatment of formalin- fixed paraffin embedded tissue sections may enhance the immunostaining in overfixed tissues. Recommended positive control: tonsil.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Factor VIII related antigen isolated from human plasma
Specificity:	This antibody reacts with factor VIII related antigen. It stains endothelial cells and megakaryocytes in various human tissues. This antibody does not cross-react with any other cell types. Cellular localization: cytoplasmic.
Formulation:	Immunoglobulin fraction containing 0.05% sodium azide as preservative State: Purified State: Liquid
Purification:	Purified
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Database Link:	Entrez Gene 7450 Human P04275
Synonyms:	vWF, von Willebrand antigen 2, von Willebrand antigen II, F8VWF, Factor VIII Related Antigen



[View online »](#)

Product images:



Formalin fixed paraffin embedded human tonsil stained with Factor VIII R antigen polyclonal rabbit antibody