

## **Product datasheet for DP044**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Gastrin (GAST) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

**Recommended Dilution:** Immunohistochemistry on formalin-fixed paraffin embedded sections:

diluted of 1:50-1:100 in an ABC method (30 minutes at room temperature)

Recommended Positive Control: Stomach

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

Immunogen: Recombinant Gastrin conjugated to KLH.

**Specificity:** The antibody is specific against gastrin-containing cells. This antibody reacts with human

gastrin I and human gastrin I fragments.

Cellular Localization: Cytoplasmic

**Formulation:** containing sodium azide as preservative

State: Purified State: Liquid serum

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Gene Name: gastrin

**Database Link:** Entrez Gene 2520 Human

P01350





## Gastrin (GAST) Rabbit Polyclonal Antibody - DP044

Background:

Gastrin is mainly known by its first described role as a gastric acid-hormone but the protein also has the ability to promote cellular growth. More recently, gastrin has been suggested to induce leukocyte-endothelial cell interactions and to have a pro-inflammatory effect. Gastrin has two biologically active peptide forms, G34 and G17 and they activate two different receptors: the CCK-1 receptor, which has low affinity for gastrin but high affinity for the related hormone cholecystokinin (CCK), and the CCK-2 receptor, which has high affinity for both gastrin and CCK and mediates the acid-secretory as well as the proliferative effects of gastrin.

Synonyms:

GAST, GAS