

Product datasheet for KN200258RB

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

B4GALT7 Human Gene Knockout Kit (CRISPR)

Product data:

Product Type: Knockout Kits (CRISPR)

Format: 2 gRNA vectors, 1 RFP-BSD donor, 1 scramble control

Donor DNA: RFP-BSD
Symbol: B4GALT7
Locus ID: 11285

Components: KN200258G1, B4GALT7 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN200258G2, B4GALT7 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002) **KN200258RBD**, donor DNA containing left and right homologous arms and RFP-BSD

functional cassette.

GE100003, scramble sequence in pCas-Guide vector

Disclaimer: These products are manufactured and supplied by OriGene under license from ERS. The kit is

designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the

experimental process.

 RefSeq:
 NM 007255

 UniProt ID:
 Q9UBV7

Synonyms: EDSP1; XGALT1; XGPT1

Summary: This gene is a member of the beta-1,4-galactosyltransferase (beta4GalT) family. Family

members encode type II membrane-bound glycoproteins that appear to have exclusive specificity for the donor substrate UDP-galactose. Each beta4GalT member has a distinct function in the biosynthesis of different glycoconjugates and saccharide structures. As type II membrane proteins, they have an N-terminal hydrophobic signal sequence that directs the

protein to the Golgi apparatus which then remains uncleaved to function as a

transmembrane anchor. The enzyme encoded by this gene attaches the first galactose in the common carbohydrate-protein linkage (GlcA-beta1,3-Gal-beta1,3-Gal-beta1,4-Xyl-beta1-O-Ser) found in proteoglycans. This enzyme differs from other beta4GalTs because it lacks the conserved Cys residues found in beta4GalT1-beta4GalT6 and it is located in cis-Golgi instead

of trans-Golgi. Mutations in this gene have been associated with the progeroid form of Ehlers-

Danlos syndrome. [provided by RefSeq, Oct 2009]





Product images:

Donor Vector Edited Chromosome



RFP, Luc, and mBFP will be under native gene promoter