

## Product datasheet for **KN200294RB**

### SKP2 Human Gene Knockout Kit (CRISPR)

#### Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 RFP-BSD donor, 1 scramble control
Donor DNA:	RFP-BSD
Symbol:	SKP2
Locus ID:	6502
Components:	<b>KN200294G1</b> , SKP2 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002) <b>KN200294G2</b> , SKP2 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002) <b>KN200294RBD</b> , donor DNA containing left and right homologous arms and RFP-BSD functional cassette. <b>GE100003</b> , scramble sequence in pCas-Guide vector
Disclaimer:	These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process.
RefSeq:	<a href="#">NM_001243120</a> , <a href="#">NM_005983</a> , <a href="#">NM_032637</a>
UniProt ID:	<a href="#">Q13309</a>
Synonyms:	FBL1; FBXL1; FLB1; p45



[View online »](#)

**Summary:**

This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbls class; in addition to an F-box, this protein contains 10 tandem leucine-rich repeats. This protein is an essential element of the cyclin A-CDK2 S-phase kinase. It specifically recognizes phosphorylated cyclin-dependent kinase inhibitor 1B (CDKN1B, also referred to as p27 or KIP1) predominantly in S phase and interacts with S-phase kinase-associated protein 1 (SKP1 or p19). In addition, this gene is established as a protooncogene causally involved in the pathogenesis of lymphomas. Alternative splicing of this gene generates three transcript variants encoding different isoforms. [provided by RefSeq, Jul 2011]

**Product images:**
