

Product datasheet for **KN200485BN**

CPT1A Human Gene Knockout Kit (CRISPR)

Product data:

| | |
|---------------|---|
| Product Type: | Knockout Kits (CRISPR) |
| Format: | 2 gRNA vectors, 1 mBFP-Neo donor, 1 scramble control |
| Donor DNA: | mBFP-Neo |
| Symbol: | CPT1A |
| Locus ID: | 1374 |
| Components: | KN200485G1 , CPT1A gRNA vector 1 in pCas-Guide CRISPR vector (GE100002) KN200485G2 , CPT1A gRNA vector 2 in pCas-Guide CRISPR vector (GE100002) KN200485BND , donor DNA containing left and right homologous arms and mBFP-Neo functional cassette. GE100003 , scramble sequence in pCas-Guide vector |
| Disclaimer: | These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process. |
| RefSeq: | NM_001031847 , NM_001876 |
| UniProt ID: | P50416 |
| Synonyms: | CPT1; CPT1-L; L-CPT1 |
| Summary: | The mitochondrial oxidation of long-chain fatty acids is initiated by the sequential action of carnitine palmitoyltransferase I (which is located in the outer membrane and is detergent-labile) and carnitine palmitoyltransferase II (which is located in the inner membrane and is detergent-stable), together with a carnitine-acylcarnitine translocase. CPT I is the key enzyme in the carnitine-dependent transport across the mitochondrial inner membrane and its deficiency results in a decreased rate of fatty acid beta-oxidation. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008] |



[View online »](#)

Product images:

