

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for KN200568LP

# Integrin beta 4 binding protein (EIF6) Human Gene Knockout Kit (CRISPR)


### **Product data:**

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 Luciferase-Puro donor, 1 scramble control
Donor DNA:	Luciferase-Puro
Symbol:	Integrin beta 4 binding protein
Locus ID:	3692
Components:	<b>KN200568G1</b> , Integrin beta 4 binding protein gRNA vector 1 in pCas-Guide CRISPR vector (GE100002) <b>KN200568G2</b> , Integrin beta 4 binding protein gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)
	<b>KN200568LPD</b> , donor DNA containing left and right homologous arms and Luciferase-Puro functional cassette. <b>GE100003</b> , scramble sequence in pCas-Guide vector
Disclaimer:	These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process.
RefSeq:	<u>NM 001267810, NM 002212, NM 181466, NM 181467, NM 181468, NM 181469, NR 052022, NR 052022, NR 052024</u>
UniProt ID:	<u>P56537</u>
Synonyms:	b(2)gcn; CAB; eIF-6; EIF3A; ITGB4BP; p27(BBP); p27BBP
Summary:	Hemidesmosomes are structures which link the basal lamina to the intermediate filament cytoskeleton. An important functional component of hemidesmosomes is the integrin beta-4 subunit (ITGB4), a protein containing two fibronectin type III domains. The protein encoded by this gene binds to the fibronectin type III domains of ITGB4 and may help link ITGB4 to the intermediate filament cytoskeleton. The encoded protein, which is insoluble and found both in the nucleus and in the cytoplasm, can function as a translation initiation factor and prevent the association of the 40S and 60S ribosomal subunits. Multiple non-protein coding transcript variants and variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Jun 2012]



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## **Product images:**



RFP, Luc, and mBFP will be under native gene promoter

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US