

Product datasheet for KN201222BN

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville. MD 20850. US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PLAUR Human Gene Knockout Kit (CRISPR)

Product data:

Product Type: Knockout Kits (CRISPR)

Format: 2 gRNA vectors, 1 mBFP-Neo donor, 1 scramble control

Donor DNA: mBFP-Neo

Symbol: PLAUR

Locus ID: 5329

Components: KN201222G1, PLAUR gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN201222G2, PLAUR gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN201222BND, donor DNA containing left and right homologous arms and mBFP-Neo

functional cassette.

GE100003, scramble sequence in pCas-Guide vector

Disclaimer: These products are manufactured and supplied by OriGene under license from ERS. The kit is

designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the

experimental process.

RefSeq: NM 001005376, NM 001005377, NM 001301037, NM 002659

UniProt ID: Q03405

Synonyms: CD87; U-PAR; UPAR; URKR

Summary: This gene encodes the receptor for urokinase plasminogen activator and, given its role in

localizing and promoting plasmin formation, likely influences many normal and pathological processes related to cell-surface plasminogen activation and localized degradation of the

extracellular matrix. It binds both the proprotein and mature forms of urokinase

plasminogen activator and permits the activation of the receptor-bound pro-enzyme by plasmin. The protein lacks transmembrane or cytoplasmic domains and may be anchored to the plasma membrane by a glycosyl-phosphatidylinositol (GPI) moiety following cleavage of the nascent polypeptide near its carboxy-terminus. However, a soluble protein is also produced in some cell types. Alternative splicing results in multiple transcript variants encoding different isoforms. The proprotein experiences several post-translational cleavage

reactions that have not yet been fully defined. [provided by RefSeq, Jul 2008]





Product images:

Donor Vector Edited Chromosome



RFP, Luc, and mBFP will be under native gene promoter