

## Product datasheet for **KN201959BN**

### **XBP1 Human Gene Knockout Kit (CRISPR)**

#### **Product data:**

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 mBFP-Neo donor, 1 scramble control
Donor DNA:	mBFP-Neo
Symbol:	XBP1
Locus ID:	7494
Components:	<b>KN201959G1</b> , XBP1 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002) <b>KN201959G2</b> , XBP1 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002) <b>KN201959BND</b> , donor DNA containing left and right homologous arms and mBFP-Neo functional cassette. <b>GE100003</b> , scramble sequence in pCas-Guide vector
RefSeq:	<a href="#">NM_001079539</a> , <a href="#">NM_005080</a>
UniProt ID:	<a href="#">P17861</a>
Synonyms:	TREB-5; TREB5; XBP-1; XBP2
Summary:	This gene encodes a transcription factor that regulates MHC class II genes by binding to a promoter element referred to as an X box. This gene product is a bZIP protein, which was also identified as a cellular transcription factor that binds to an enhancer in the promoter of the T cell leukemia virus type 1 promoter. It may increase expression of viral proteins by acting as the DNA binding partner of a viral transactivator. It has been found that upon accumulation of unfolded proteins in the endoplasmic reticulum (ER), the mRNA of this gene is processed to an active form by an unconventional splicing mechanism that is mediated by the endonuclease inositol-requiring enzyme 1 (IRE1). The resulting loss of 26 nt from the spliced mRNA causes a frame-shift and an isoform XBP1(S), which is the functionally active transcription factor. The isoform encoded by the unspliced mRNA, XBP1(U), is constitutively expressed, and thought to function as a negative feedback regulator of XBP1(S), which shuts off transcription of target genes during the recovery phase of ER stress. A pseudogene of XBP1 has been identified and localized to chromosome 5. [provided by RefSeq, Jul 2008]



[View online »](#)

## Product images:

