

## Product datasheet for KN202069LP

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **CXCR4 Human Gene Knockout Kit (CRISPR)**

#### **Product data:**

**Product Type:** Knockout Kits (CRISPR)

**Format:** 2 gRNA vectors, 1 Luciferase-Puro donor, 1 scramble control

**Donor DNA:** Luciferase-Puro

Symbol: CXCR4 Locus ID: 7852

**Components:** KN202069G1, CXCR4 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN202069G2, CXCR4 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN202069LPD, donor DNA containing left and right homologous arms and Luciferase-Puro

functional cassette.

**GE100003**, scramble sequence in pCas-Guide vector

**Disclaimer:** These products are manufactured and supplied by OriGene under license from ERS. The kit is

designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the

experimental process.

RefSeq: NM 001008540, NM 003467, NM 001348056, NM 001348059, NM 001348060

UniProt ID: <u>P61073</u>

Synonyms: CD184; D2S201E; FB22; HM89; HSY3RR; LAP-3; LAP3; LCR1; LESTR; NPYR; NPYR; NPYRL;

NPYY3R; WHIM

Summary: This gene encodes a CXC chemokine receptor specific for stromal cell-derived factor-1. The

protein has 7 transmembrane regions and is located on the cell surface. It acts with the CD4 protein to support HIV entry into cells and is also highly expressed in breast cancer cells. Mutations in this gene have been associated with WHIM (warts, hypogammaglobulinemia, infections, and myelokathexis) syndrome. Alternate transcriptional splice variants, encoding

different isoforms, have been characterized. [provided by RefSeq, Jul 2008]





# **Product images:**

### Donor Vector Edited Chromosome



RFP, Luc, and mBFP will be under native gene promoter