

# **Product datasheet for KN202079RB**

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## IL1 beta (IL1B) Human Gene Knockout Kit (CRISPR)

**Product data:** 

**Product Type:** Knockout Kits (CRISPR)

**Format:** 2 gRNA vectors, 1 RFP-BSD donor, 1 scramble control

Donor DNA: RFP-BSD Symbol: IL1 beta Locus ID: 3553

**Components: KN202079G1**, IL1 beta gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

**KN202079G2**, IL1 beta gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN202079RBD, donor DNA containing left and right homologous arms and RFP-BSD

functional cassette.

**GE100003**, scramble sequence in pCas-Guide vector

**Disclaimer:** These products are manufactured and supplied by OriGene under license from ERS. The kit is

designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the

experimental process.

**RefSeq:** <u>NM 000576</u>

UniProt ID: P01584

Synonyms: IL-1; IL1-BETA; IL1F2




#### **Summary:**

The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. Similarly, IL-1B has been implicated in human osteoarthritis pathogenesis. Patients with severe Coronavirus Disease 2019 (COVID-19) present elevated levels of pro-inflammatory cytokines such as IL-1B in bronchial alveolar lavage fluid samples. The lung damage induced by the Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is to a large extent, a result of the inflammatory response promoted by cytokines such as IL-1B. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. [provided by RefSeq, Jul 2020]

## **Product images:**

### Donor Vector Edited Chromosome



RFP, Luc, and mBFP will be under native gene promoter