

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for KN202718RB

Alpha B Crystallin (CRYAB) Human Gene Knockout Kit (CRISPR)

Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 RFP-BSD donor, 1 scramble control
Donor DNA:	RFP-BSD
Symbol:	Alpha B Crystallin
Locus ID:	1410
Components:	 KN202718G1, Alpha B Crystallin gRNA vector 1 in pCas-Guide CRISPR vector (GE100002) KN202718G2, Alpha B Crystallin gRNA vector 2 in pCas-Guide CRISPR vector (GE100002) KN202718RBD, donor DNA containing left and right homologous arms and RFP-BSD functional cassette. GE100003, scramble sequence in pCas-Guide vector
Disclaimer:	These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process.
RefSeq:	<u>NM 001289807, NM 001289808, NM 001330379, NM 001885, NM 001368245, NM 001368245</u>
UniProt ID:	<u>P02511</u>
Synonyms:	CMD1II; CRYA2; CTPP2; CTRCT16; HEL-S-101; HSPB5; MFM2




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Scrigene Alpha B Crystallin (CRYAB) Human Gene Knockout Kit (CRISPR) – KN202718RB

Mammalian lens crystallins are divided into alpha, beta, and gamma families. Alpha crystallins Summary: are composed of two gene products: alpha-A and alpha-B, for acidic and basic, respectively. Alpha crystallins can be induced by heat shock and are members of the small heat shock protein (HSP20) family. They act as molecular chaperones although they do not renature proteins and release them in the fashion of a true chaperone; instead they hold them in large soluble aggregates. These heterogeneous aggregates consist of 30-40 subunits; the alpha-A and alpha-B subunits have a 3:1 ratio, respectively. Two additional functions of alpha crystallins are an autokinase activity and participation in the intracellular architecture. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. Alpha-A and alpha-B gene products are differentially expressed; alpha-A is preferentially restricted to the lens and alpha-B is expressed widely in many tissues and organs. Elevated expression of alpha-B crystallin occurs in many neurological diseases; a missense mutation cosegregated in a family with a desmin-related myopathy. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2019]

Product images:



RFP, Luc, and mBFP will be under native gene promoter

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US