

Product datasheet for KN202883BN

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Ube1L (UBA7) Human Gene Knockout Kit (CRISPR)

Product data:

Product Type: Knockout Kits (CRISPR)

Format: 2 gRNA vectors, 1 mBFP-Neo donor, 1 scramble control

Donor DNA: mBFP-Neo

Symbol: Ube1L Locus ID: 7318

Components: KN202883G1, Ube1L gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN202883G2, Ube1L gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN202883BND, donor DNA containing left and right homologous arms and mBFP-Neo

functional cassette.

GE100003, scramble sequence in pCas-Guide vector

Disclaimer: These products are manufactured and supplied by OriGene under license from ERS. The kit is

designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the

experimental process.

RefSeq: <u>NM 003335</u>

UniProt ID: P41226

Synonyms: D8; UBA1B; UBE1L; UBE2

Summary: The modification of proteins with ubiquitin is an important cellular mechanism for targeting

abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E1 ubiquitin-activating enzyme family. The encoded enzyme is a retinoid target that triggers promyelocytic leukemia (PML)/retinoic acid receptor alpha (RARalpha) degradation and apoptosis in acute

promyelocytic leukemia, where it is involved in the conjugation of the ubiquitin-like


interferon-stimulated gene 15 protein. [provided by RefSeq, Jul 2008]





Product images:

Donor Vector Edited Chromosome



RFP, Luc, and mBFP will be under native gene promoter