

Product datasheet for KN202896RB

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cystatin SN (CST1) Human Gene Knockout Kit (CRISPR)

Product data:

Product Type: Knockout Kits (CRISPR)

Format: 2 gRNA vectors, 1 RFP-BSD donor, 1 scramble control

Donor DNA: RFP-BSD
Symbol: Cystatin SN

Locus ID: 1469

Components: KN202896G1, Cystatin SN gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN202896G2, Cystatin SN gRNA vector 2 in pCas-Guide CRISPR vector (GE100002) **KN202896RBD**, donor DNA containing left and right homologous arms and RFP-BSD

functional cassette.

GE100003, scramble sequence in pCas-Guide vector

Disclaimer: These products are manufactured and supplied by OriGene under license from ERS. The kit is

designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the

experimental process.

P01037

RefSeq: <u>NM 001898</u>

UniProt ID:

Synonyms: cystatin 1; cystatin SA-I; cystatin SN; cysteine proteinase inhibitor, type 2 family;

OTTHUMP00000030444: OTTHUMP00000164184

Summary: The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences.

Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin

locus on chromosome 20 contains the majority of the type 2 cystatin genes and

pseudogenes. This gene is located in the cystatin locus and encodes a cysteine proteinase inhibitor found in saliva, tears, urine, and seminal fluid. [provided by RefSeq, Jul 2008]





Product images:

Donor Vector Edited Chromosome



RFP, Luc, and mBFP will be under native gene promoter