

## Product datasheet for KN204878RB

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **Glucocorticoid Receptor (NR3C1) Human Gene Knockout Kit (CRISPR)**

**Product data:** 

**Product Type:** Knockout Kits (CRISPR)

**Format:** 2 gRNA vectors, 1 RFP-BSD donor, 1 scramble control

**Donor DNA:** RFP-BSD

Symbol: Glucocorticoid Receptor

**Locus ID:** 2908

**Components: KN204878G1**, Glucocorticoid Receptor gRNA vector 1 in pCas-Guide CRISPR vector

(GE100002)

KN204878G2, Glucocorticoid Receptor gRNA vector 2 in pCas-Guide CRISPR vector

(GE100002)

KN204878RBD, donor DNA containing left and right homologous arms and RFP-BSD

functional cassette.

GE100003, scramble sequence in pCas-Guide vector

RefSeq: NM 000176, NM 001018074, NM 001018075, NM 001018076, NM 001018077,

NM 001020825, NM 001024094, NM 001204258, NM 001204259, NM 001204260, NM 001204261, NM 001204262, NM 001204263, NM 001204264, NM 001364182, NM 001364184, NR 157096, NM 001364180, NM 001364181,

NM 001364183, NM 001364185

UniProt ID: P04150

Synonyms: GCCR; GCR; GCRST; GR; GRL

**Summary:** This gene encodes glucocorticoid receptor, which can function both as a transcription factor

that binds to glucocorticoid response elements in the promoters of glucocorticoid responsive genes to activate their transcription, and as a regulator of other transcription factors. This receptor is typically found in the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in

target tissues. Mutations in this gene are associated with generalized glucocorticoid

resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities

(PMID:15866175). [provided by RefSeq, Feb 2011]





# **Product images:**

#### Donor Vector Edited Chromosome



RFP, Luc, and mBFP will be under native gene promoter