

Product datasheet for KN205288BN

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MCSF Receptor (CSF1R) Human Gene Knockout Kit (CRISPR)

Product data:

Product Type: Knockout Kits (CRISPR)

Format: 2 gRNA vectors, 1 mBFP-Neo donor, 1 scramble control

Donor DNA: mBFP-Neo

Symbol: MCSF Receptor

Locus ID: 1436

Components: KN205288G1, MCSF Receptor gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN205288G2, MCSF Receptor gRNA vector 2 in pCas-Guide CRISPR vector (GE100002) **KN205288BND**, donor DNA containing left and right homologous arms and mBFP-Neo

functional cassette.

GE100003, scramble sequence in pCas-Guide vector

Disclaimer: These products are manufactured and supplied by OriGene under license from ERS. The kit is

designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the

experimental process.

RefSeq: NM 001288705, NM 005211, NR 109969, NM 001349736

UniProt ID: <u>P07333</u>

Synonyms: C-FMS; CD115; CSF-1R; CSFR; FIM2; FMS; HDLS; M-CSF-R

Summary: The protein encoded by this gene is the receptor for colony stimulating factor 1, a cytokine

which controls the production, differentiation, and function of macrophages. This receptor mediates most if not all of the biological effects of this cytokine. Ligand binding activates the receptor kinase through a process of oligomerization and transphosphorylation. The

receptor kinase through a process of oligomerization and transphosphorylation. The encoded protein is a tyrosine kinase transmembrane receptor and member of the CSF1/PDGF receptor family of tyrosine-protein kinases. Mutations in this gene have been

associated with a predisposition to myeloid malignancy. The first intron of this gene contains a transcriptionally inactive ribosomal protein L7 processed pseudogene oriented in the opposite direction. Alternative splicing results in multiple transcript variants. Expression of a splice variant from an LTR promoter has been found in Hodgkin lymphoma (HL), HL cell lines

and anaplastic large cell lymphoma. [provided by RefSeq, Mar 2017]





Product images:

Donor Vector Edited Chromosome



RFP, Luc, and mBFP will be under native gene promoter