

Product datasheet for **KN206042RB**

TRAF6 Human Gene Knockout Kit (CRISPR)

Product data:

Product Type: Knockout Kits (CRISPR)

Format: 2 gRNA vectors, 1 RFP-BSD donor, 1 scramble control

Donor DNA: RFP-BSD

Symbol: TRAF6

Locus ID: 7189

Components: **KN206042G1**, TRAF6 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)
KN206042G2, TRAF6 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)
KN206042RBD, donor DNA containing left and right homologous arms and RFP-BSD functional cassette.
GE100003, scramble sequence in pCas-Guide vector

Disclaimer: These products are manufactured and supplied by OriGene under license from ERS. The kit is designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the experimental process.

RefSeq: [NM_004620](#), [NM_145803](#)

UniProt ID: [Q9Y4K3](#)

Synonyms: MGC:3310; RNF85



[View online »](#)

Summary:

The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins are associated with, and mediate signal transduction from, members of the TNF receptor superfamily. This protein mediates signaling from members of the TNF receptor superfamily as well as the Toll/IL-1 family. Signals from receptors such as CD40, TNFSF11/RANCE and IL-1 have been shown to be mediated by this protein. This protein also interacts with various protein kinases including IRAK1/IRAK, SRC and PKCzeta, which provides a link between distinct signaling pathways. This protein functions as a signal transducer in the NF-kappaB pathway that activates I kappa B kinase (IKK) in response to proinflammatory cytokines. The interaction of this protein with UBE2N/UBC13, and UBE2V1/UEV1A, which are ubiquitin conjugating enzymes catalyzing the formation of polyubiquitin chains, has been found to be required for IKK activation by this protein. This protein also interacts with the transforming growth factor (TGF) beta receptor complex and is required for Smad-independent activation of the JNK and p38 kinases. This protein has an amino terminal RING domain which is followed by four zinc-finger motifs, a central coiled-coil region and a highly conserved carboxyl terminal domain, known as the TRAF-C domain. Two alternatively spliced transcript variants, encoding an identical protein, have been reported. [provided by RefSeq, Feb 2012]

Product images: