

## Product datasheet for **KN208513BN**

### FAM20B Human Gene Knockout Kit (CRISPR)

#### Product data:

Product Type:	Knockout Kits (CRISPR)
Format:	2 gRNA vectors, 1 mBFP-Neo donor, 1 scramble control
Donor DNA:	mBFP-Neo
Symbol:	FAM20B
Locus ID:	9917
Components:	<b>KN208513G1</b> , FAM20B gRNA vector 1 in pCas-Guide CRISPR vector (GE100002) <b>KN208513G2</b> , FAM20B gRNA vector 2 in pCas-Guide CRISPR vector (GE100002) <b>KN208513BND</b> , donor DNA containing left and right homologous arms and mBFP-Neo functional cassette. <b>GE100003</b> , scramble sequence in pCas-Guide vector
RefSeq:	<a href="#">NM_014864</a> , <a href="#">NM_001324310</a> , <a href="#">NM_001324311</a>
UniProt ID:	<a href="#">O75063</a>
Synonyms:	gxl1
Summary:	Responsible for the 2-O-phosphorylation of xylose in the glycosaminoglycan-protein linkage region of proteoglycans thereby regulating the amount of mature GAG chains. Sulfated glycosaminoglycans (GAGs), including heparan sulfate and chondroitin sulfate, are synthesized on the so-called common GAG-protein linkage region (GlcUA $\beta$ 1-3Gal $\beta$ 1-3Gal $\beta$ 1-4Xyl $\beta$ 1-O-Ser) of core proteins, which is formed by the stepwise addition of monosaccharide residues by the respective specific glycosyltransferases. Xylose 2-O-phosphorylation may influence the catalytic activity of B3GAT3 (GlcAT-I) which completes the precursor tetrasaccharide of GAG-protein linkage regions on which the repeating disaccharide region is synthesized.[UniProtKB/Swiss-Prot Function]



[View online »](#)

## Product images:

